

Phytotoxic, Cytotoxic and Insecticidal Activities of *Chrysophthalmum dichotomum* Boiss. and Heldr.

Fatma Ayaz¹, Nurgün Küçükboyacı^{2*}, Barış Bani³, Bilge Şener², Muhammad Iqbal Choudhary⁴

¹Department of Pharmacognosy, Faculty of Pharmacy, Selcuk University, Konya, TURKEY.

²Department of Pharmacognosy, Faculty of Pharmacy, Gazi University, Ankara, TURKEY.

³Department of Biology, Faculty of Arts and Science, Kastamonu University, Kastamonu, TURKEY.

⁴International Center for Chemical Sciences, University of Karachi, Karachi, PAKISTAN.

ABSTRACT

Background: *Chrysophthalmum dichotomum* Boiss. and Heldr. (Asteraceae) is an endemic herbaceous plant to Southern part of Turkey. **Aim:** To investigate *in vitro* phytotoxic, cytotoxic and insecticidal activities of *C. dichotomum*. **Methods:** The MeOH extract of *C. dichotomum* was fractionated through subsequent solvent extractions in increasing polarity with *n*-hexane, chloroform and *n*-butanol. The MeOH extract and its fractions were evaluated for their biological activities using *in vitro* screening bioassays such as cytotoxicity on brine shrimps, phytotoxicity against *Lemna minor* and insecticidal activity against *Rhyzopertha dominica* and *Tribolium castaneum*. **Results:** The *n*-hexane and chloroform fractions showed significant phytotoxic activity (100 % growth inhibition) at 1000 µg/ml against *L. minor*. The brine shrimp lethality test revealed that the chloroform and remaining water fractions of *C. dichotomum* have moderate and positive lethality with LD₅₀ values of 169.48 and 46.26 µg/mL, respectively. In addition, the chloroform and *n*-butanol fractions had low and moderate insecticidal activity with 20 and 40 % of mortality against *Tribolium castaneum*, respectively. **Conclusion:** This study demonstrates that *C. dichotomum* consists of bioactive constituents responsible for phytotoxicity, cytotoxicity on brine shrimps and insecticidal activity.

Key words: *Chrysophthalmum dichotomum*, Asteraceae, Phytotoxicity against *Lemna minor*, Cytotoxicity on brine shrimps, Insecticidal activity.

INTRODUCTION

The genus *Chrysophthalmum* Schultz Bip. (Asteraceae, tribus: Inulaeae) and consists of four species all over the world. In Turkey, the genus is represented by three species, namely *C. montanum* (DC.) Boiss., *C. dichotomum* Boiss. and Heldr. and *C. gueneri* Aytac and Anderb. *C. dichotomum* is an endemic herbaceous plant that grows in wooded or shrubby valley beds in the district of Antalya, Turkey.¹ Up to date, no phytochemical and biological data has been reported on the plant. In our previous study, we firstly evaluated the cytotoxicity of *C. dichotomum* against selected cancer cell lines by Sulforhodamine B (SRB) assay.²

As a part of our continuing researches on the genus *Chrysophthalmum*, we recently evaluated the crude methanol extract and its

fractions of *C. montanum* for their *in vitro* phytotoxic, cytotoxic and insecticidal activities. In that study, the crude extract, *n*-hexane and chloroform fractions have positive lethality on brine shrimp as well as *n*-hexane and chloroform fractions had significant phytotoxicity (100% of growth inhibition) at 1000 µg/mL.³ In other our previous study, we found that chloroform extract of *C. montanum* and its isolated sesquiterpene lactones have significant cytotoxic activities on different cancer cell lines using SRB assay.⁴ Family Asteraceae is characterized by structurally diverse sesquiterpenes and, therefore it has been extensively studied in point of cytotoxic and anticancer activities. Sesquiterpenes have also been reported to use as toxic or insect-feeding deterrent.⁵⁻⁶

Submission Date: 20-10-2017;

Revision Date: 05-12-2017;

Accepted Date: 17-05-2018

DOI: 10.5530/ijper.52.4s.105

Correspondence:

Prof. Nurgün Küçükboyacı,

Department of

Pharmacognosy, Faculty of

Pharmacy, Gazi University,

06330 Ankara, TURKEY.

E-mail: nurgun@gazi.edu.tr



www.ijper.org

Our ongoing researches on the genus *Chrysophthalmum*, we now aimed to investigate *in vitro* phytotoxic, cytotoxic and insecticidal activities of *C. dichotomum*.

MATERIALS AND METHODS

Plant material

Chrysophthalmum dichotomum were collected from damp places in opening of *Pinus nigra* forest around Tinaztepe cave, in Konya in August, 2014. The plant was identified by one of our authors Barış Bani PhD, (Kastamonu University). Voucher specimen (F. Ayaz 46) was deposited at Herbarium of Gazi University (GAZI), Ankara, Turkey.

Preparation of extracts

The air-dried whole plants of *C. dichotomum* (150 g) were extracted four times (4×1500 mL) with 80% methanol at 25°C by stirring for 2 days. Following filtration, the combined methanol extracts were evaporated *in vacuo* at 40°C to dryness. The concentrated MeOH extract (25.8 g for CD) were further fractionated by successive solvent extractions with *n*-hexane (11×200 mL), chloroform (8×200 mL) and *n*-butanol saturated with H₂O (7×200 mL) in a separatory funnel. Each extract and remaining water phase (R-H₂O) after solvent extractions were evaporated to dryness under reduced pressure to yield “*n*-Hexane fraction” (1.1 g CDH), “CHCl₃ fraction” (2.6 g for CDC), “*n*-BuOH fraction” (7.9 g for CDB) and “R-H₂O fraction” (11.6 g CDR), respectively.

Phytochemical analysis

The CHCl₃ extract of *C. gueneri* (1 mg/mL) was applied to silica gel plate. After developing with the mixture of chloroform: acetone (80:20) as mobile phases, TLC plate was evaluated under UV light at 254 and 366 nm for the determination of fluorescent compounds. Anisaldehyde reagent was sprayed on the plates to visualize the separated compounds and then plates were heated for 5 min at 100°C. Sesquiterpenes appeared with pink and purple coloration.

Phytotoxicity against *Lemna minor*

The *Lemna minor* L. phytotoxicity assay was performed for the extracts of *C. dichotomum* by adopting the protocol as described by Atta-ur-Rahman.⁷ The medium was prepared by mixing various constituents in 1000 mL distilled water and pH was adjusted 6.0-7.0 by adding KOH pellets. The extracts (30.0 mg) were dissolved in 1.5 mL of methanol (stock solution). The stock solutions of the extracts were diluted to get final concentrations as 10, 100 and 1000 µg/mL. The solvent was allowed to evaporate overnight under sterile condition. To each

flask was then added 20 mL medium and 10 plants, each one containing a rosette of two fronds of *L. minor*. Other flasks were supplemented with medium and reference plant growth inhibitor (Paraquate) as negative and positive controls, respectively. All flasks were kept in growth cabinet for 7 days. The number of fronds per flasks was counted and recorded at the end of the incubation period. The growth regulation as a percentage (%) was determined using the formula given below:

$$\text{Growth regulation (\%)} = \frac{100 - \text{Number of fronds in test samples}}{\text{Number of fronds in negative control}} \times 100$$

The activity criteria indicate that the growth regulation (%) of 0-39% for low activity, 40-59% for moderate activity, 60-69% for good activity and >70% for significant activity were detected.

Cytotoxicity on brine shrimps

Brine shrimp (*Artemia saline* Leach) eggs (50 mg) were sprinkled in a hatching tank (a rectangular dish 22x32 cm) half-filled with filtered brine solution. The extracts (20 mg) were dissolved in 2 mL of methanol (stock solution). The concentrations of 10, 100 and 1000 µg/mL of each extracts from stock solutions were prepared in three vials and the solvent was evaporated by keeping over night. 2 days of hatching later, 30 shrimps were added in each vial with the volume adjusted to 5 ml using sea water. The vials were incubated at 25-27°C for 24 h under illumination. Other vials were supplemented with solvent, and reference cytotoxic drug (Etoposide: 7.4625 µg/ml) which served as – ve and + ve controls, respectively. The number of brine shrimps that survived was counted in each vial and LD₅₀ values with 95% confidence intervals were determined using Finney computer software.⁸⁻⁹

Insecticidal activity

The extracts of *C. dichotomum* were tested against *Rhyzopertha dominica* and *Tribolium castaneum* using impregnated filter paper method.¹⁰ The extracts (200 mg) were dissolved in 3 mL of methanol (stock solution). The extracts were applied to filter paper (1019.10 µg/cm²) of appropriate size (9 cm or 90 mm) on petri plates using micropipette. The plates were left for 24 h to evaporate the solvent. The next day, 10 insects of each species were placed in each plate [test and permethrin (239.5 µg/cm²) was used as + ve control; methanol was used as – ve control] using a clean brush. The plates were incubated at 27°C for 24 h with 50% relative humidity in

the growth chamber. For the calculation, the number of survivals of each species was counted and mortality (%) was determined using the following formula:

$$\text{Mortality (\%)} = \frac{100 - \text{Number of insects alive in test samples}}{\text{Number of insects alive in control}} \times 100$$

RESULTS

In our study, we investigated the methanol extract and its fractions of *C. dichotomum* for their *in vitro* phytotoxic, cytotoxic and insecticidal activities. The phytotoxicity of the tested samples on *L. minor* was observed to have dose dependent activity, because low activity was detected in the chloroform and *n*-butanol fractions with 19.0 and 10.6% inhibition at 100 µg/mL, respectively, as well as in remaining water fraction with 37.5% inhibition at 1000 µg/mL. Moderate phytotoxic activity was found in the *n*-butanol fraction (42.6% inhibition) at 1000 µg/mL. Significant phytotoxic activity was shown in the *n*-hexane and chloroform fractions of the plant; 100.0% inhibition for each fraction at 1000 µg/mL (Table 1).

The cytotoxic properties of the extract and fractions of *C. dichotomum* were investigated at a concentration of 10, 100 and 1000 µg/mL, using etoposide as a standard. The chloroform and remaining water fractions had moderate and positive lethality with LD₅₀ values of 169.48 and 46.26 µg/mL against the brine shrimps, respectively (Table 2).

The extract and fractions of *C. dichotomum* were also screened for their insecticidal effects against *Rhyzopertha dominica* and *Tribolium castaneum* using permethrin as a standard drug. The chloroform and *n*-butanol fractions had low and moderate insecticidal activities with 20 and 40% of mortality against *Tribo-*

Table 1: Phytotoxic activities of the extract and fractions of *C. dichotomum*.

Sample	% Growth inhibition		
	10 µg/mL	100 µg/mL	1000 µg/mL
CD	0	0	0
CDH	0	0	100.0
CDC	0	19.0	100.0
CDB	0	10.6	42.6
CDR	0	0	37.5

Standard drug: Paraquate (0.015 µg/mL)

Table 2: Cytotoxic activities of the extract and fractions of *C. dichotomum*.

Sample	No of survivals from 30 shrimps			LD ₅₀ (µg/mL)
	10 µg/mL	100 µg/mL	1000 µg/mL	
CD	23	22	17	5900.97
CDH	28	25	21	-
CDC	24	25	03	169.48
CDB	18	16	15	771.45
CDR	16	15	12	46.26

Standard drug: Etoposide (LD₅₀ =7.46 µg/mL)

Table 3: Insecticidal activities of the extract and fractions of *C. dichotomum*.

Sample	% Mortality	
	<i>Tribolium castaneum</i>	<i>Rhyzopertha dominica</i>
CD	0	0
CDH	0	0
CDC	20	0
CDB	40	0
CDR	0	0

Reference insecticide: Permethrin (239.5 µg/cm²)

lium castaneum, respectively. There was no insecticidal effect on all tested samples against *Rhyzopertha dominica* (Table 3).

DISCUSSION

To date, a number of plant extracts prepared in different solvents was screened to determine the preliminary cytotoxic activity on brine shrimp larvae. Brine shrimp lethality assay is a preliminary method for the assessment of cytotoxic compounds which could show a significant correlation with their anticancer potentials.¹¹⁻¹³ In addition, to develop natural herbicides and insecticides that are safe and user friendly for the environment is important for screening in terms of phytotoxic and insecticidal activities of plants.¹⁴

According to our results, the chloroform fraction of *C. dichotomum* was found as a promising sample due to having cytotoxicity on brine shrimps. In our previous study, the chloroform fraction of *C. dichotomum* exhibited cytotoxicity on MCF-7, MDA-MB-231, LNCaP, PC-3 (lung) and HT-29 cancer cell lines using SRB assay with % viability values 22.99, 21.08, 24.27, 16.02 and 28.00, respectively.² Our preliminary phytochemical analysis of the bioactive chloroform fraction by TLC depicted that sesquiterpenes were detected as principal components.

CONCLUSION

The present study firstly showed that the potential of *C. dichotomum* on biological activities such as cytotoxicity on brine shrimps, phytotoxic and insecticidal effects *in vitro*. Further investigations are continuing on the isolation and characterization for the responsible bioactive constituents from *C. dichotomum*.

ACKNOWLEDGEMENT

This study was supported by TUBİTAK-2214/A and ICCBS-HEJ. TUBİTAK-2214/A, TUBİTAK-2211/A and ICCBS-HEJ.

CONFLICT OF INTEREST

The authors declare no conflict of interest..


ABBREVIATIONS USED

SRB: Sulforhodamine B; **CD:** The Concentrated MeOH extract; **CDH:** n-Hexane fraction; **CDC:** CHCl₃ Fraction; **CDB:** n-BuOH fraction; **CDR:** R-H₂O Fraction.

REFERENCES

- Grierson AJC. *Chrysophthalmum* Schultz Bip. In: Flora of Turkey and the East Aegean Islands, Davis, PH, ed., Vol 5, Edinburgh: Edinburgh University Press. 1975;5:52-3.
- Ayaz F, Kucukboyacı N, Sarimahmut M, Bani B, Duman H, Ulukaya E, et al. "Cytotoxic Activity of the Genus *Chrysophthalmum* Schultz Bip. From Turkey Against Various Human Cancer Cell Lines", 3rd EACR-Sponsored Anticancer Agent Development Congress. 2015;42. 2015, İzmir, Turkey.
- Ayaz F, Küçükboyacı N, Duman H, Şener B, Choudhary MI. Cytotoxic, Phytotoxic and Insecticidal Activities of *Chrysophthalmum montanum* (DC.) Boiss Turk. J Pharm Sci. 2017;14:290-3.
- Ayaz F, Kucukboyacı N, Goren N, Calis I, Aydinlik S, Ulukaya E, et al. "Bioassay-Guided Isolation and Cytotoxic Effects of Extract and Chemical Constituents of *Chrysophthalmum montanum* (DC.) Boiss. Of Turkish Origin", Turk. J Biochem. 2016;41(S4):46.
- Fraga BM. Natural sesquiterpenoids. Nat Prod Rep. 2004;21:669-93.
- Rodriguez E, Towers GH, Mitchell JC. Biological activities of sesquiterpene lactones. Phytochemistry. 1976;15(11):1573-80.
- Atta-ur-Rahman. Studies in Natural Product Chemistry. The Netherlands: Elsevier Science Publishers BV. 1991;9:383-409.
- Meyer BN, Ferrigni NR, Putnam JE, Jacobsen LB, Nicholas PE, McLaughlin JL. Brine Shrimp: A convenient general bioassay for active plant constituents. Planta Med. 1982;45(5):31-4.
- Finny DJ. Probit Analysis. 3rd ed. Cambridge: Cambridge University Press. 1971:333.
- Atta-ur-Rahman, Choudhary MI, William JT. Bioassay techniques for drug development. The Netherlands: Harward Academic Publisher. 2001:67-8.
- Ullah R, Ibrar M, Shah S, Hameed I. Phytotoxic, cytotoxic and insecticidal activities of *Calendula arvensis* L. E3 J Biotechnol Pharm. Res. 2012;3(6):104-11.
- Khuda F, Iqbal Z, Khan A, Nasir F, Muhammad N, Khan JA, et al. Metal analysis, phytotoxic, insecticidal and cytotoxic activities of selected medicinal plants of Khyber Pakhtunkhwa. Pak J Pharm. Sci. 2012;25(1):51-8.
- Uddin G, Rehman TU, Arfan M, Liaqat W, Mohammad G, Choudhary MI. Phytochemical analysis, antifungal and phytotoxic activity of seeds of medicinal plant *Indigofera heterantha* Wall. Middle-East J Sci Res. 2011;8:603-5.
- Saeed M, Muhammad N, Khan H, Khan SA. Analysis of toxic heavy metals in branded Pakistani Herbal Products. J Chem Soc Pak. 2010;32(4):471-5.

PICTORIAL ABSTRACT



SUMMARY

- In the present study was investigated on *C. dichotomum* on biological activities such as cytotoxicity
- brine shrimps, cytotoxicity and insecticidal effects *in vitro*.
- The chloroform fraction of *C. dichotomum* showed-moderate cytotoxicity on brine shrimps.
- Significant phytotoxic activity was shown in the n-hexane and chloroform fractions of the plant.

About Authors



Dr. Ayaz: Has been graduated from Hacettepe Univ., Fac. of Pharmacy in 2011 as the Second Degree. She has completed her Ph.D. (Pharmacognosy) with "Pharmacognosic Researches on *Chrysophthalmum* Schultz Bip. Species Growing in Turkey" in 2017. In 2018, she became an Assist. Professor in Selcuk Univ. Fac. of Pharmacy. She has been working in the Department of Pharmacognosy, Faculty of Pharmacy, Selcuk University, Konya, Turkey.



Prof. Küçükboyacı: Received her PhD degree in Pharmacognosy in 1999 from Gazi University, Ankara, Turkey. She is currently the Head of the Department of Phytopharmacy, Faculty of Pharmacy, Gazi University. Her research interests include the development of natural compounds from medicinal plants and the evaluation of their biological activities as well as phytomedicines. She has been working in the Department of Pharmacognosy, Faculty of Pharmacy, Gazi University, Ankara, Turkey.



Dr. Baris Bani: Is working in the Department of Biology from Kastamonu University, Kastamonu, Turkey. His research area is plant taxonomy.



Prof. Sener: Has been graduated from Ankara Univ., Fac. of Pharmacy in 1974 as the First Degree. She has completed her Ph.D. (Pharmacognosy) awarded by TUBITAK in 1977. In 1981, she became an Assoc. Professor. She was involved within the process of establishing Dept. of Pharmacognosy, Fac. of Pharmacy, Gazi University in 1982. She achieved some researches at the Dept. of Chemistry, The Pennsylvania State University (USA) 1986-1988 as Visiting scientist awarded by NSF. She became a full Professor in 1988. Prof. Sener also involved several administrative works at Gazi University; as Director at the Dept. of Pharmacognosy, Chair of the Division, Co-Director at the Inst. of Health Sciences and Dean at the Faculty of Pharmacy. Her main research field is development of bioactive compounds, evaluation of Phytomedicines and Nutraceuticals. She has won 21 awards. The degree of "Adjunct Professor" was given to her by University of Karachi, Pakistan. She has completed 48 projects from NATO, NSF, IUPAC, Soctrates-Grundtvig, TUBITAK and University Research Funds. 11 books, 88 chapters and 662 research articles alongwith 68 conferences and 315 plenary, invited lectures for her credit. She has supervised 5 Ph.D., 13 M.Sc. at Gazi University as well as co-supervised 17 Ph.D. thesis at the University of Karachi. She is a member for 14 scientific societies.



Dr. Muhammad Iqbal Choudhary Is Director and Professor of Bioorganic and Natural Product Chemistry at the International Center for Chemical and Biological Sciences (H. E. J. Research Institute of Chemistry and Dr. Panjwani Center for Molecular Medicine and Drug Research). Prof. Choudhary has, since 1990, been among the world leaders in the field of natural product chemistry, and has made pioneering contributions in the discovery of novel natural products. He has discovered many potent anti-epileptic and anti-leishmanial compounds from indigenous medicinal plants that are under clinical trials. His contributions to reverse bacterial resistance to antibiotics represent seminal contributions in this important field. He leads the developing world's finest research center of natural product chemistry (HEJ Research Institute of Chemistry) since 2002, and has trained hundreds of young researchers, especially women, from across the Afro-Asian region in natural product chemistry. He has established several research centers in Pakistan, and helped to setup research units in Africa, and South and Central Asia. His scientific, and capacity building contributions have been recognized by prestigious national and international awards and honors, and fellowships of several academies of science. Prof. Choudhary has 1016 publications (Citations 13,000, h index 44) in the fields of organic and bioorganic chemistry, along with 60 international patents (46 US Patents), 59 books and 40 chapters in books, published by major U.S. and European presses. On the basis of his researches, 73 students have been awarded Ph.D. degrees in various areas of natural product and bioorganic chemistry.

Cite this article: Ayaz F, Küçükboyacı N, Bani B, Sener B, Choudhary MI. Phytotoxic, Cytotoxic and Insecticidal Activities of *Chrysophthalmum dichotomum* Boiss. and Heldr. Indian J of Pharmaceutical Education and Research. 2018;529(4 Suppl 2):s252-s256.