

# Therapeutic Effect of Hydro-Ethanollic Extract of *Pothos scandens* L on key Carbohydrate Metabolizing Enzymes and Xenobiotic Marker Enzymes in DMBA Induced Experimental Mammary Carcinoma

Jethinlalkhosh Jayapalan Prasannakumari<sup>1\*</sup>, Praveena Padmam<sup>2</sup>, Victor Arokia Doss<sup>3</sup>

<sup>1</sup>Research and Development Centre, Bharathiar University, Coimbatore -641046, Tamil Nadu, INDIA.

<sup>2</sup>Department of Biotechnology Engineering, Sahrdaya College of Engineering and Technology, Thrissur, Kerala, INDIA.

<sup>3</sup>Department of Biochemistry, P S G College of Arts and Science, Coimbatore - 641014, Tamil Nadu, INDIA.

## ABSTRACT

**Objective:** In the present study, phytochemical evaluation and the therapeutic effect of 50 % hydro-ethanollic extract of aerial parts of *Pothos scandens* (HEEPS) on glycolytic enzymes, TCA cycle enzymes, gluconeogenic enzymes and xenobiotic marker enzymes in the liver and breast tissues of 7, 12 -Dimethylbenz [a] anthracene (DMBA) induced breast cancer bearing rats were carried out. **Materials and Methods:** Five groups of (each group six rats) female albino Spargue-Dawley rats were taken. Group I (control), Group II (DMBA induced, 20 mg/ kg b.wt p.o), Group III (DMBA + Cyclophosphamide, 10mg/kg b.wt p.o), Group IV (DMBA + 200 mg/kg b.wt p.o of HEEPS) and Group V (DMBA + 400 mg/kg b.wt p.o of HEEPS). The alteration in the level of carbohydrate metabolizing enzymes such as Hexokinase, Phosphogluco isomerase, Glucose-6-phosphatase, Fructose-1,6-bis phosphatase, Succinate dehydrogenase and Malate dehydrogenase together with the level of xenobiotic markers like Cytochrome P450 reductase, Cytochrome C oxidase and Glutathione -S- transferase were analyzed in all the groups. GC-MS analysis of the plant extract was also carried out. **Results:** Increased level of glycolytic enzymes, decreased level of TCA cycle enzymes, gluconeogenic enzymes and altered level of xenobiotic marker enzymes observed in Group II was brought to near normal in treatment groups in a dose dependent manner suggesting potential anticancer activity of HEEPS. This was supported by the GC-MS analysis of the plant extract which revealed the presence of many compounds with anticancer activity. **Conclusion:** The study revealed that HEEPS is effective in restoring the activity of mitochondria in DMBA induced breast cancer.

**Key words:** *Pothos scandens* L, GC-MS, 7, 12 -Dimethylbenz [a] anthracene (DMBA), Mammary Carcinoma, Carbohydrate Metabolizing Enzymes, Xenobiotic Markers.

Submission Date: 18-01-2017;

Revision Date: 14-03-2017;

Accepted Date: 18-03-2017

DOI: 10.5530/ijper.51.3.70

### Correspondence:

Jethinlalkhosh J P, Quality Control Section, Food Corporation of India, District Office, Palakkad - 678731, Kerala, INDIA.  
Phone no: +919633860543  
E mail: mailjethin@gmail.com

## INTRODUCTION

Cancer can be described as a group of more than 100 different diseases which are characterized by uncontrolled cellular growth, local tissue invasion and distant metastasis.<sup>1</sup> Cancer claims over 6 million lives every year and it continues to represent the largest cause of mortality in the world.<sup>2</sup> Breast cancer is the most common form of cancer and the leading cause of cancer mortality among women worldwide.<sup>3,4</sup> The number of new breast cancer cases in India is

about 100,000 per year.<sup>5</sup> It is a major clinical problem that possesses significant social and economic challenges to the health care system.<sup>6</sup> Breast cancer is a collection of breast diseases that have diverse genomic variations histopathologies and clinical outcomes.<sup>7</sup>

Mitochondrial dysfunction, rapid growth and cellular proliferation are hallmarks of tumor cells. In order to support continuous cell growth and proliferation in a hypoxic



www.ijper.org

and acidic conditions generated during progressive tumor cell growth, cancer cells alter their metabolism by shifting the burden of energy metabolism from mitochondrial oxidative phosphorylation to glycolysis.<sup>8</sup> Molecular defects in mitochondrial or nuclear genes that code for proteins that are involved in mitochondria biogenesis; respiratory maintenance of membrane potential; chain assembly and function and energy metabolic or signal transducing pathways can be the cause of mitochondrial dysfunction.<sup>9</sup> Despite various therapeutic strategies including chemotherapy to treat cancer, drug resistance and high systemic toxicity limit the successful outcomes in most cases. Moreover cancer treatment is usually accompanied by diverse side effects to different body organs.<sup>10</sup> Hence there is a global trend to go back to natural resources (medicinal plants) which are more therapeutically effective, culturally acceptable and economically affordable.<sup>11</sup> Phytochemicals with anticancer potential which interfere with carcinogenesis makes them interesting tools in cancer research.<sup>12</sup>

*Pothos scandens* (*P. scandens*), a medicinal aroid, is a climbing shrub with aerial roots growing on trees and rocks like ivy which belongs to the family Araceae. After being fried in oil, the bruised root of the plant is applied to promote healing of abscesses. An infusion of the leaves of this plant is used by the Indian people as a bath for curing convulsions and epilepsy. Apart from that, the stem is also widely used to treat asthma. It has been also reported that the whole plant is used against various health problems and disorders such as diarrhea<sup>13</sup>, small pox<sup>14</sup>, muscle catches, sprains<sup>15</sup> and bone fracture.<sup>16</sup> Recently, the ethanolic extract of *P. scandens* has been reported to be effective in wound healing<sup>17</sup> and the plant has also been reported to have significant antioxidant and antipyretic activity.<sup>18</sup> The ethanolic extracts of aerial part of *P. scandens* showed the presence of phytochemicals such as alkaloids, glycosides, flavonoids and phenolic compounds and has been shown to inhibit mast cell derived immediate-type allergic reactions and mast cell degranulation.<sup>19</sup> The methanolic extract of leaves of *P. scandens* has been reported to have cytotoxic as well as thrombolytic potential.<sup>20</sup> From the literature survey it was found that the phytochemical analysis of leaf extract of *P. scandens* showed the presence of alkaloid, catechin, coumarin, flavonoid, phenolics, saponins, sugar, glycoside, xanthoprotein<sup>21</sup> and phytosterols.<sup>18</sup> 1,2- Benzenedicarboxylic acid, diisooctylester; n-Hexadecanoic acid, 9,12,15-Octadecatrienoic acid, (ZZZ)-; Octadecanoic acid; Phytol and 9, 12 Octadecadienoic acid (ZZ) were detected as major compounds in the GC-MS analysis of the ethanolic extract of *P. scandens* leaf.<sup>22</sup>

As per the ethnobotanical data collected during the field surveys made on several visits between 2004 and 2006 to three Akha communities in Chiang Rai in northern Thailand, it was found that the whole aerial parts of *P. scandens* were used by the traditional healers to treat cancer.<sup>23</sup> However, on thorough literature survey, it was found that there were no studies related to the anticancer properties of this plant. Hence, the present study aims to investigate the therapeutic properties of *P. scandens* for its potential anticancer activity by evaluating their effect on carbohydrate metabolizing enzymes and biotransformation enzymes in DMBA induced experimental mammary carcinoma.

## MATERIALS AND METHODS

### Chemicals and reagents

7, 12 -Dimethylbenz [a] anthracene (DMBA) was obtained from Sigma Chemicals Co. (St. Louis, MO, USA). All other chemicals and solvents used were of analytical grade.

### Plant material

*P. scandens* were collected from in and around Palai, Kottayam, Kerala. They were identified and certified by the Taxonomist, Botanical Survey of India (BSI), Coimbatore, Tamil Nadu, India (Plant identification no.-BSI/SRC/5/23/2013-14/Tech/685).

### Extraction Procedure

The aerial parts of *P. scandens* were shade dried and ground to coarse powder. The coarse powder was extracted using 50% ethanol using soxhlet apparatus for 48 hours. The extracts were filtered and condensed to dryness using a rotary evaporator and stored at -20°C for further studies.

### Animals

Female albino Spargue-Dawley rats (150-200 g) used in the present study was procured from the animal breeding station, Mannuthy, Kerala, India. They were housed in polypropylene cages (38 × 23 × 10cm) with not more than six animals per cage and maintained under standard environmental conditions (14h dark /10h light cycles; temp 25±2°C; 35-60% humidity, air ventilation) and were fed with standard pellet diet (M/s. Hindustan Lever Ltd., Mumbai, India) and fresh water *ad libitum*. The animals were acclimatized to the environment for two weeks prior to experimental use. Animals were fasted over night before the experimental schedule, but have free access for water *ad libitum*. Ethical clearance for the handling of experimental animals was obtained from the Institutional Animal Ethics Committee

(IAEC) constituted for the purpose and care of laboratory animals as per the guidance of the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), Ministry of Social Justice and Empowerment, Government of India (IAEC No. ML-EA-CPCSEA/09-2014/02).

### Experimental Design

The animals were randomly divided into five groups of six animals each. Group I served as control rats. Group II rats were induced with a single dose of DMBA (20 mg/kg b. wt, dissolved in 0.75 ml sunflower oil and 0.25 ml physiological saline) by oral gavage at the first day of the experimental period. Group III received Cyclophosphamide (CP) (10mg/kg p.o), Group IV received 200 mg/kg b.wt and Group V received 400 mg/kg b.wt of HEEPS orally for 60 days after 90 days of DMBA induction. At the end of the experimental period, all the animals were anesthetized with diethyl ether, and they were sacrificed by decapitation. Animals were starved overnight before sacrifice. The breast and liver tissues were excised and rinsed 2 to 3 times in ice cold normal saline followed by 0.15 M Tris- HCl (pH 7.4), blotted dried and known weight of breast and liver were homogenized in 0.25 M Tris-HCl buffer (pH 7.5). The homogenate was subjected to differential centrifugation at 4°C. The cell organelle such as mitochondria, microsomes and cytosolic fractions were isolated. Total homogenate and subcellular fractions were used for the assay of the following parameters in breast and liver samples.

### Biochemical Analysis

The following biochemical parameters in the liver and breast tissue were analysed. Hexokinase (HK), Branstrup *et al.*,<sup>24</sup>; Phosphogluco isomerase (PGI), Horrocks *et al.*,<sup>25</sup>; Glucose-6-phosphatase (G-6-Pase), King<sup>26</sup>; Fructose-1, 6-bis phosphatase (F-1,6-BPase), Gancedo and Gancedo<sup>27</sup>; Succinate dehydrogenase (SDH), Slater and Bonner<sup>28</sup>; Malate dehydrogenase (MDH), Mehler *et al.*,<sup>29</sup>; Cytochrome P450 reductase (POR), Masters *et al.*,<sup>30</sup>; Cytochrome C oxidase (COX), Wharton and Tzagoloff<sup>31</sup> and Glutathione -S- transferase (GST) Habig *et al.*,<sup>32</sup>

### Gas Chromatography-Mass Spectrometry (GC-MS) analysis of hydro-ethanolic extract of *P. scandens* (HEEPS)

GC- MS analysis of HEEPS was performed with Shimadzu GC- MS (model QP2010) operating in EI mode at 70 eV with a Restek -5MS column (30m × 0.25mm × 0.25µm). Helium gas was used as carrier gas at a constant flow rate 1ml/ min and an injection volume was 1µL. The oven temperature programs was initially kept at 150°C and hold for 1 min, and then with a rise

of 10°C/ minute upto 250°C and hold for 3 min, and again with a rise of 10°C/ min upto 280°C and hold for 5 min. Injector temperature 260°C; ion source temperature 250°C, quadrupole temperature 150°C, interface temperature 250°C, full scan mode, scan range 40-600 m/z and solvent delay of 4 min was employed.

The interpretation of the spectrum from GC- MS analysis was done using the database of National Institute of Standard and Technology (NIST). By comparing the mass spectrum of the unknown component with the known component in the NIST library; the name, structure and molecular weight of the components of the test materials were ascertained.

### Statistical analysis

The results are expressed as mean ± standard deviation (SD). Differences between groups were assessed by one-way analysis of variance (ANOVA) using the Statistical Package of Social Sciences (SPSS, Version 16.0 for windows). The group means were compared by Duncan's Multiple Range Test (DMRT). Values were considered statistically significant when  $p < 0.05$ .

## RESULTS

### Effect of HEEPS on carbohydrate metabolizing enzymes

The effect of HEEPS on carbohydrate metabolizing enzymes in the liver and mammary gland tissue of mammary carcinoma bearing rats is given in Table 1. The result indicated that the activity of glycolytic enzymes namely HK and PGI in the liver and mammary tissues of rats bearing mammary carcinoma (Group II) was significantly increased by 81.82% and 84.13% in liver and by 60% and 67.74% in mammary tissue respectively when compared with control rats (Group I). However, it was seen that the activity reverted to near normal when treated with plant extract (Group IV and Group V). The activity shown by the treatment group was almost near to those treated with standard drug (Group III).

The activity of TCA cycle enzymes like SDH and MDH were significantly lowered by 75.22% and 76.09% in liver and by 71.85% and 72.18% in mammary tissue respectively in Group II rats compared to the Group I rats. It was observed from the result that the treatment groups (Group III, Group IV and Group V) showed near normal activity of these enzymes.

Gluconeogenic enzymes like G-6-Pase and F-1, 6-BPase showed a decrease in activity by 52.86% and 51.60% in liver and by 68.72% and 48.39% in mammary tissue respectively in Group II rats when compared with control rats whereas the near normal activity was shown



Decreased activity of gluconeogenic enzymes viz G-6-Pase and F-1, 6-BPase might lead to an increase in the concentration of their substrates glucose-6-phosphate and fructose-1, 6-bisphosphate. The resultant accumulated glucose-6-phosphate might be utilized for the pentose phosphate pathway to produce ribose-5-phosphate for the rapid synthesis of nucleotide in cancer bearing cells. On the other hand fructose-1, 6-bisphosphate acts as an allosteric activator of pyruvate kinase.<sup>43</sup> The activities of these enzymes were returned to near normal when treated with plant extract suggesting a protective role of plant.

POR is a flavoenzyme ubiquitously expressed in all cells including malignant cells and catalyzes the electron transfer required for the P450 enzyme activity.<sup>44</sup> It has been confirmed from studies that POR overexpression substantially increases the potency of prodrug mediated tumor cell killing.<sup>45-48</sup> In the present study, a decrease in the activity of POR in both liver and breast tissues has been observed in cancer induced rats (Group II) which was brought to near normal in groups treated with plant extract. This suggests the capability of plant extract in augmenting the expression of POR.

COX is a protein present in the inner membrane of mitochondria and is responsible for the conversion of

**Table 1: Effect of HEEPS on carbohydrate metabolizing enzymes of liver and breast tissue against DMBA induced tumor**

Parameter	Group I (Control)	Group II (DMBA)	Group III (DMBA+CP)	Group IV (DMBA + HEEPS 200 mg/kgbw)	Group V (DMBA + HEEPS 400 mg/kgbw)
LIVER					
HK	0.011 ± 0.002	0.020 ± 0.003*	0.011 ± 0.001*	0.014 ± 0.002*	0.013 ± 0.001*
PGI	0.063 ± 0.010	0.116 ± 0.022*	0.062 ± 0.005*	0.076 ± 0.011*	0.070 ± 0.011*
SDH	1.703 ± 0.18	0.422 ± 0.08*	1.325 ± 0.21*	0.672 ± 0.05*	1.040 ± 0.07*
MDH	1.748 ± 0.13	0.418 ± 0.03*	1.263 ± 0.32*	0.623 ± 0.07*	1.098 ± 0.05*
G-6-Pase	0.367 ± 0.06	0.173 ± 0.04*	0.344 ± 0.03*	0.289 ± 0.04*	0.314 ± 0.01*
F-1, 6-BPase	0.281 ± 0.05	0.136 ± 0.03*	0.264 ± 0.02*	0.200 ± 0.02*	0.243 ± 0.02*
BREAST					
HK	0.045 ± 0.006	0.072 ± 0.012*	0.047 ± 0.004*	0.051 ± 0.004*	0.049 ± 0.003*
PGI	0.040 ± 0.018	0.124 ± 0.020*	0.041 ± 0.017*	0.085 ± 0.017*	0.052 ± 0.020*
SDH	1.197 ± 0.18	0.337 ± 0.03*	1.042 ± 0.17*	0.602 ± 0.05*	0.933 ± 0.08*
MDH	6.568 ± 0.96	1.827 ± 0.51*	5.938 ± 0.89*	3.642 ± 0.70*	5.068 ± 0.43*
G-6-Pase	0.227 ± 0.01	0.071 ± 0.04*	0.194 ± 0.01*	0.147 ± 0.03*	0.176 ± 0.02*
F-1, 6-BPase	0.746 ± 0.07	0.385 ± 0.10*	0.688 ± 0.08*	0.586 ± 0.04*	0.639 ± 0.04*

Values are expressed as mean ± SD for six rats in each group. \*p<0.05. Group I vs Group II, Group II vs Group III, Group IV and Group V. Units: HK- μ moles of glucose 6 phosphate formed/min/mg protein; PGI - μ moles of fructose formed/min/mg protein; SDH - μmoles of succinate oxidized /min/mg protein; MDH - μmoles of NADH oxidized/min/mg protein; G-6-Pase and F-1, 6- BPase - μ moles of phosphorous liberated/min/mg protein

**Table 2: Effect of HEEPS on xenobiotic marker enzymes of liver and breast tissue against DMBA induced tumor**

Parameter	Group I (Control)	Group II (DMBA)	Group III (DMBA+CP)	Group IV (DMBA + HEEPS 200 mg/kgbw)	Group V (DMBA + HEEPS 400 mg/kgbw)
LIVER					
POR	20.038 ± 0.33	8.795 ± 0.58*	15.780 ± 0.28*	11.612 ± 0.31*	13.098 ± 0.52*
COX	223.143 ± 1.18	49.810 ± 5.39*	160.650 ± 1.63*	135.670 ± 6.15*	151.075 ± 7.01*
GST	52.800 ± 1.29	22.118 ± 1.71*	43.463 ± 1.24*	36.803 ± 1.15*	39.868 ± 1.94*
BREAST					
POR	11.680 ± 1.54	3.605 ± 0.89*	9.587 ± 1.48*	6.750 ± 1.59*	7.817 ± 1.07*
COX	475.923 ± 8.57	161.510 ± 8.81*	412.637 ± 6.48*	292.653 ± 6.37*	372.192 ± 7.27*
GST	43.742 ± 5.27	21.300 ± 5.86*	37.872 ± 6.18*	31.813 ± 2.32*	35.908 ± 4.19*

Values are expressed as mean ± SD for six rats in each group. \*p<0.05. Group I vs Group II, Group II vs Group III, Group IV and Group V. Units: POR - μ moles of Cytochrome reduced/min/mg protein; COX - μ moles of Cytochrome oxidised/min/mg protein; GST - μ moles of CDNB conjugation formed/min/mg protein

**Table 3: Phytochemical compounds identified through the GC-MS study of HEEPS**

Sl. No	Retention Time (min)	Compound	Molecular Formula	Molecular Weight
1	9.940	3,7- Dimethylnonane	C <sub>11</sub> H <sub>24</sub>	156
2	12.553	Silane, trimethyl (1-methyl dodecyl oxy)-	C <sub>16</sub> H <sub>36</sub> OSi	272
3	14.552	1,3-Di isopropoxy-1,3-dimethyl-1,3-disilacyclobutane	C <sub>10</sub> H <sub>24</sub> O <sub>2</sub> Si <sub>2</sub>	232
4	15.935	Sym-tetra (isopropyl) disiloxane	C <sub>12</sub> H <sub>30</sub> OSi <sub>2</sub>	246
5	16.569	α Curcumene	C <sub>15</sub> H <sub>22</sub>	202
6	17.318	α Bergamotene	C <sub>15</sub> H <sub>24</sub>	204
7	22.830	Butyldimethyl (2-styryl[1,3]dithian-2-yl) silane	C <sub>18</sub> H <sub>28</sub> S <sub>2</sub> Si <sub>2</sub>	336
8	23.575	6-Methyl-6-(5-methylfuran-2-yl) heptan-2-one	C <sub>13</sub> H <sub>20</sub> O <sub>2</sub>	208
9	24.556	1,1,3,3-Tetramethylcyclopentane	C <sub>9</sub> H <sub>18</sub>	126
10	25.760	5-Isopropenyl-2-methyl-7-oxabicyclo[4.1.0]heptan-2-ol	C <sub>10</sub> H <sub>16</sub> O <sub>2</sub>	168
11	26.954	3,7,11,15-Tetramethyl-2-hexadecen-1-ol (Synonym: Phytol)	C <sub>20</sub> H <sub>40</sub> O	296
12	27.083	Hexahydrofarnesyl acetone	C <sub>18</sub> H <sub>36</sub> O	268
13	27.425	3,7,11,15-Tetramethyl-2-hexadecen-1-ol (Synonym: Phytol)	C <sub>20</sub> H <sub>40</sub> O	296
14	27.749	3,7,11,15-Tetramethyl-2-hexadecen-1-ol (Synonym: Phytol)	C <sub>20</sub> H <sub>40</sub> O	296
15	28.337	Hexahydrofarnesyl acetone	C <sub>18</sub> H <sub>36</sub> O	268
16	28.519	Hexadecanoic acid, methyl ester	C <sub>17</sub> H <sub>34</sub> O <sub>2</sub>	270
17	28.725	2-Octenoic acid, 4-isopropylidene-7-methyl-6-methylene-, methyl ester	C <sub>14</sub> H <sub>22</sub> O <sub>2</sub>	222
18	28.875	1-Hexadecen-3-ol, 3,7,11,15-tetramethyl- (Synonym: Isophytol)	C <sub>20</sub> H <sub>40</sub> O	296
19	29.585	Ethyl heptadecanoate	C <sub>19</sub> H <sub>38</sub> O <sub>2</sub>	298
20	31.021	9,12-Octadecadienoic acid (Z,Z)-, methyl ester	C <sub>19</sub> H <sub>34</sub> O <sub>2</sub>	294
21	31.090	9-Octadecenoic acid (Z)-, methyl ester	C <sub>19</sub> H <sub>36</sub> O <sub>2</sub>	296
22	31.158	9-Octadecenoic acid (Z)-, methyl ester	C <sub>19</sub> H <sub>36</sub> O <sub>2</sub>	296
23	31.260	3,7,11,15-Tetramethyl-2-hexadecen-1-ol (Synonym: Phytol)	C <sub>20</sub> H <sub>40</sub> O	296
24	31.416	Octadecanoic acid, methyl ester	C <sub>19</sub> H <sub>38</sub> O <sub>2</sub>	298
25	32.256	Ethyl octadecanoate	C <sub>20</sub> H <sub>40</sub> O <sub>2</sub>	312
26	33.469	Eicosane	C <sub>20</sub> H <sub>42</sub>	282
27	34.216	5-methyl-5-(4,8,12-trimethyltridecyl)oxolan-2-one	C <sub>21</sub> H <sub>40</sub> O <sub>2</sub>	324
28	34.675	Dotriacontane	C <sub>32</sub> H <sub>66</sub>	450
29	35.128	Nopol	C <sub>11</sub> H <sub>18</sub> O	166
30	36.093	Hexatriacontane	C <sub>36</sub> H <sub>74</sub>	506
31	37.021	Mono (2-ethylhexyl)phthalate	C <sub>16</sub> H <sub>22</sub> O <sub>4</sub>	278
32	37.830	Tetrapentacontane	C <sub>54</sub> H <sub>110</sub>	758
33	40.005	Tetrapentacontane	C <sub>54</sub> H <sub>110</sub>	758
34	41.807	Methyl steviol	C <sub>21</sub> H <sub>32</sub> O <sub>3</sub>	332
35	42.527	1-Bromotriacontane	C <sub>30</sub> H <sub>61</sub> Br	501
36	43.190	Docosa-2,6,10,14,18-pentaen-22-al, 2,6,10,15,18-pentamethyl-, all trans	C <sub>27</sub> H <sub>44</sub> O	384
37	44.416	Stigmast-5-en-3-ol, (3 beta) (Synonym: β Sitosterol)	C <sub>29</sub> H <sub>50</sub> O	414
38	45.783	5-Nitrobenzofuran-2-one	C <sub>8</sub> H <sub>5</sub> NO <sub>4</sub>	179
39	46.823	Cedrol	C <sub>15</sub> H <sub>26</sub> O	222

molecular oxygen to water in the terminal complex of mitochondrial respiratory chain.<sup>49</sup> In the present study reduction in the activity of the enzyme was found in DMBA induced breast cancer. Defective complex IV has been observed in various studies conducted in many human tumors and mouse xenografts.<sup>50</sup> The activity of the enzyme was significantly increased in the treatment groups indicating the capability of *P. scandens* extract to repair the mitochondrial membrane damage. Similar results were observed in the studies conducted using tangeretin in breast cancer bearing rats.<sup>51</sup>

GST is a major group of phase II enzymes for which glutathione is the substrate. The expression of GST is controlled by multiple transcription factors and is highly inducible by dietary components.<sup>52</sup> The enzyme is involved in the detoxification of variety of xenobiotics and also plays an important role in carcinogen detoxification.<sup>53,54</sup> In the present study, the level of GST was decreased in carcinoma bearing rats depicting the deleterious effect of carcinogen. *P. scandens* extract treated groups showed near normal activity showing curative effect of the plant extract. Chemopreventive compounds act as blocking agents that affect the initiation phase of cancer by detoxification of carcinogen through increasing the activity of GST.<sup>55</sup> Our results are in accordance with the above findings as we have observed a significant decline in the activity of GST in DMBA induced cancer bearing animals (Group II) and a significant increase in the activity of the enzyme in treatment groups.

The HEEPS contains rich phytochemicals which was identified by GC-MS analysis. In the present study in terms of percentage phytol,  $\beta$ -sitosterol, hexadecanoic acid methyl ester, octadecanoic acid methyl esters were predominant in the extract. Phytol has been reported to have anticancer<sup>56</sup>, antimicrobial<sup>57</sup>, diuretic and anti-inflammatory activity.<sup>58</sup>  $\beta$ -sitosterol has been reported to impart protection to cells against damage by reactive oxygen species by increasing the activities of antioxidant enzymes.<sup>59</sup> Also it was pointed out in the study of Awad *et al.*,<sup>60</sup> that  $\beta$ -sitosterol can alleviate cancer development by reducing production of carcinogens and its treatment has reduced the breast cancer cell growth.<sup>61</sup> Hexadecanoic acid methyl ester was reported to have anticancer and antioxidant properties and octadecanoic acid has been reported to have anticancer, antioxidant antibacterial, antifungal, hypocholesterolemic, antihistamic and anti-coronary properties.<sup>62</sup> Apart from these, remaining compounds mentioned in Table 3 detected by GC-MS has been reported to have significant application in drug industry.

## CONCLUSION

Damage to the mitochondrial membrane and increased rate of glycolytic enzymes during cancer is an important field of research in breast cancer studies. From the present study, it was evident that significant damage was caused to mitochondria in breast cancer induced animals which resulted in an increase in the glycolytic enzymes and a decrease in the level of TCA cycle, gluconeogenic enzymes. Also it was found that the levels of marker enzymes such as POR, COX and GST were also altered in breast cancer induced animals. All the altered enzyme activities were brought to near normal activity in plant extract treated groups which suggest the potential therapeutic effect of the plant. The GC-MS analysis of the plant also well supported the fact that the plant contained important compound which could be employed for the treatment of breast cancer. To the best of our knowledge, this is the first publication about the therapeutic activity of *P. scandens* extract against breast cancer. However, compounds responsible for its anticancer property and its mode of action need further study.

## CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest.

## ABBREVIATION USED

**HEEPS:** 50 % hydro-ethanolic extract of *Pothos scandens*; **DMBA:** 7, 12 -Dimethylbenz [a] anthracene; **CP:** Cyclophosphamide; **HK:** Hexokinase; **PGI:** Phosphogluco isomerase; **SDH:** Succinate dehydrogenase; **MDH:** Malate dehydrogenase; **G-6-Pase:** Glucose-6-phosphatase; **F-1,6-BPase:** Fructose-1,6-bisphosphatase; **POR:** Cytochrome P450 reductase; **COX:** Cytochrome C oxidase; **GST:** Glutathione -S- transferase; **NIST:** National Institute of Standard and Technology; **SPSS:** Statistical Package of Social Sciences; **OXPHOS:** Oxidative phosphorylation.

## REFERENCES

1. Kaufman D and Chabner BA, Clinical strategies for cancer treatment: The Role of drugs. In: Cancer chemotherapy and biotherapy: Principle and practice, Chabner, B.A. and D.L. Longo (Ed.). Lippincott-Raven, Philadelphia. 1996;1-16. PMID:10155984.
2. Karthikeyan R, Karthigayan S, Sri Balasubashi M, Vijayalakshni S and Balasubramanian T, Antitumor effect of snake venom (*Hydrophis spiralis*) on ehrlich ascites carcinoma bearing mice. *Int. J. Cancer Res.* 2007;s3:167-73.
3. Smigal C, Jemal A, Ward E, Cokkinides V, Smith R., Howe H, *et al.*, Trends in breast cancer by race and ethnicity: Update. *CA Cancer J Clin.* 2006;56:106-30. <https://doi.org/10.3322/canjclin.56.3.168>; PMID:16514137.
4. Russo J, Yang X and Hu Y, Biological and molecular basis of human breast cancer. *Frontiers Biosci.* 1998;3:944-60. <https://doi.org/10.2741/A335>.

5. Yadav P and Jaroli DP, Breast cancer: awareness and risk factors in college-going younger age group women in Rajasthan. *Asian Pac J Cancer Prev*. 2010;11(2):319-22. PMID:20843108.
6. Guo J and Wang MH, Extract of *Ulmus davidiana* planch barks induced apoptosis in human hepatoma cell line HEPG2. *Exp Clin Sci J*. 2009;8:130-7.
7. Vargo-Gogola T and Rosen JM, Modelling breast cancer: one size does not fit all. *Nat Rev Cancer*. 2007;7(9):659-72. <https://doi.org/10.1038/nrc2193>; PMID:17721431.
8. Warburg O, On the origin of cancer cells. *Science*. 1956;123(3191):309-14. <https://doi.org/10.1126/science.123.3191.309>.
9. Ma Y, Bai RK, Trieu R and Wong LJ, Mitochondrial dysfunction in human breast cancer cells and their transmembrane cybrids. *Biochimica et Biophysica Acta*. 2010;1797(1):29-37. <https://doi.org/10.1016/j.bbabi.2009.07.008> PMID:19647716 PMID:PMC2787670.
10. Saalu LC, Ajayi GO and Adeneye AA, Ethanolic seed extracts of grapefruit (*Citrus paradise* Macfad) on an effective attenuator of Doxorubicin-induced oxidative stress in the rat heart. *Int. J. Cancer Res*. 2009;5(2):44-52. <https://doi.org/10.3923/ijcr.2009.44.52>.
11. Jeyachandran R, Mahesh A and Cindrella L, DEN- Induced cancer and its alleviation by *Anisomeles malabarica* (L.) R.Br. ethanolic leaf extract in male albino mice. *Int. J. Cancer Res*. 2007;3:174-9. <https://doi.org/10.3923/ijcr.2007.174.179>.
12. Cai Y, Luo Q, Sun M, *et al.*, Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sci*. 2004;74(17):2157-84. <https://doi.org/10.1016/j.lfs.2003.09.047>; PMID:14969719.
13. Sangwoo L, Chunjie X and Shengji P, Ethnobotanical survey of medicinal plants at periodic markets of honghe prefecture in Yunnan Province, SW China. *J Ethnopharmacol*. 2008;117(2):362-77. <https://doi.org/10.1016/j.jep.2008.02.001>; PMID:18359178
14. Baharuddin S and Mansor M, Medicinal aroids conservation: A case study of floral garden School of Biological Scie Ces, Universiti Sains Malaysia. Proceedings of the 4th IMT-GT UNINET Conference. 2002; 216-9.
15. Bhandary MJ, Chandrashekar KR and Kaveriappa KM, Medical ethnobotany of the Siddis of Uttara Kannada district, Karnataka, India. *J Ethnopharmacol*. 1995;47(3):149-58. [https://doi.org/10.1016/0378-8741\(95\)01274-H](https://doi.org/10.1016/0378-8741(95)01274-H).
16. Das AK, Dutta BK and Sharma GD, Medicinal plants used by different tribes of Cachar district, Assam. *Indian J Tradit Knowl*. 2008;7:446-54.
17. Sainuddin T and Mohammed HKP, Formulation & pharmacological evaluation of herbal gel of *Pothos Scandens* Linn. *WebmedCentral WOUND HEALING*. 2010; 1:WMC001344.
18. Sajeesh T, Arunachalam K and Parimelazhagan T, Antioxidant and antipyretic studies on *Pothos scandens* L. *Asian Pacific J Tropical Med*, 2011; 4: 889-99. [https://doi.org/10.1016/S1995-7645\(11\)60214-9](https://doi.org/10.1016/S1995-7645(11)60214-9).
19. Gupta S, Duraiswamy B and Satishkumar MN, Peritoneal mast cell stabilization potential of *Pothos scandens* L. *Indian J.Pharmacol*. 2013;45(1):83-6. <https://doi.org/10.4103/0253-7613.106442>; PMID:23542883 PMID:PMC3608302
20. Yusuf ATM, Rashid MMU, Hassan MM, Sayeed MA and Jainul MA, *In-vitro* Cytotoxic and thrombolytic potential of *Pothos scandens* L. *J. Pharmacogn. Phytochem*. 2013;2(1):193-8.
21. Lalitharani, Mohan VR, Maruthupandian, Pharmacognostical and phytochemical studies on *Pothos scandens* L. *Int J Phytomed*, 2010;2:277-83.
22. Lalitharani S, Mohan VR, Regini GS and Kalidass C, GC-MS analysis of ethanol extract of *Pothos scandens* leaf. *J Herb Med Toxicol*, 2009;3:159-60.
23. Angkhana I, Shengji P, Balslev H, Wong PW and Trisonthi C, A comparative study on medicinal plants used in Akha's traditional medicine in China and Thailand, cultural coherence or ecological divergence? *J Ethnopharmacol*. 2008;116:508-17. <https://doi.org/10.1016/j.jep.2007.12.015>; PMID:18280071
24. Branstrup N, Krik JE and Bruni C, The hexokinase and phosphoglucoserase activities of aorta and pulmonary artery tissue in individuals of various ages. *J Gerontol*. 1957;12(2):166-71. <https://doi.org/10.1093/geronj/12.2.166>.
25. Horrocks JE, Ward J and Kind J, A routine method for the determination of phosphoglucosomerase activity in body fluid. *J Clin Pathol*. 1963;16(3):248-51. <https://doi.org/10.1136/jcp.16.3.248>.
26. King J, The hydrolases-acid and alkaline phosphatases. In: *Practical clinical Enzymology*, Van, D (Ed.), Norstand company Limited, London. 1965b;191-208.
27. Gancedo JM and Gancedo C, Fructose-1,6-bisphosphatase, Phosphofructokinase and Glucose-6-phosphodehydrogenase. *Proc.SOC.EXP.Biol. Med*. 1971;106:607-9.
28. Slater EC and Bonner WD, The effect of fluoride on the succinic oxidase system. *Biochem J*. 1952; 52(2):185-96. <https://doi.org/10.1042/bj0520185>.
29. Mehler AH, Kornberg A, Grisolia S and Ochoa S, The enzymatic mechanism of oxidation- reductions between malate or isocitrate or pyruvate. *J Biol Chem*. 1948;174:s961-77.
30. Masters BS, Williams CH and Kamin H, The preparation and properties of microsomal TPNH cytochrome c reductase from pig liver. In: Estabrook, R.W., Pullman, M.E. eds: *Methods Enzymol*. 1967;10:565-73. [https://doi.org/10.1016/0076-6879\(67\)10098-0](https://doi.org/10.1016/0076-6879(67)10098-0).
31. Wharton DC and Tzagoloff A, Cytochrome oxidase from beef heart mitochondria. *Methods Enzymol*. 1967;10:245-50. [https://doi.org/10.1016/0076-6879\(67\)10048-7](https://doi.org/10.1016/0076-6879(67)10048-7).
32. Habig WH and Jakoby WB, Glutathione S-transferases. The first enzymatic step in mercapturic acid formation. *J. Bio. Chem*. 1974;249(22):7130-9.
33. Heimann R., Heuson JC and Coune A, Tumors developing in oophorectomized Sprague-Dawley rats after a single gastric instillation of 7,12-dimethylbenz(a)anthracene. *Cancer Res*. 1968;28(2):309-13. PMID:5641521.
34. Bhuvaneshwari V, Velmurugan B, Abraham SK and Nagini S, Tomato and garlic by gavage modulate 7, 12-dimethylbenz[a]anthracene-induced genotoxicity and oxidative stress in mice. *Braz J Med Biol Res*. 2004; 37(7):1029-34. <https://doi.org/10.1590/S0100-879X2004000700012>.
35. Esterbauer H, Estimation of peroxidative damage. *Pathol Biol (Paris)*. 1996;44:25-8.
36. Yeung SJ, Pan J and Lee MH, Roles of p53, MYC and HIF-1 in regulating glycolysis: the seventh hallmark of cancer. *Cell Mol Life Sci*. 2008;65(24):3981-99. <https://doi.org/10.1007/s00018-008-8224-x>; PMID:18766298.
37. DeBerardinis R.J, Lum JJ, Hatzivassiliou G and Thompson CB, The biology of cancer:metabolic reprogramming fuels cell growth and proliferation. *Cell Metab*. 2008 7(1):11-20. <https://doi.org/10.1016/j.cmet.2007.10.002>; PMID:18177721.
38. DeBerardinis R.J, Sayed N, Ditsworth D and Thompson CB, Brick by brick: metabolism and tumor cell growth. *Curr. Opin. Genet. Dev*. 2008b;18(1):54-61. <https://doi.org/10.1016/j.gde.2008.02.003> PMID:18387799 PMID:PMC2476215
39. Parry DM and Pedersen PL, Intracellular localization and properties of particulate hexokinase in the Novikoff ascites tumour. Evidence for an outer mitochondrial membrane location. *J Biol Chem*. 1983;258:10904-12. PMID:6885806
40. Parveen V, Sharma DC and Shastri KD. A clinicobiochemical study of phosphohexose-isomerase, alioesterate and sulphhydryl group in 25 normal and 35 cancer patients. *Indian J cancer*. 1974;43:448-56.
41. Dimitrova-Shumkovska J, Veenman L, Ristoski T, *et al.*, Decreases in binding capacity of the mitochondrial 18 kda translocator protein accompany oxidative stress and pathological signs in rat liver after DMBA exposure. *Toxicol Pathol*. 2010;38(6):957-68. <https://doi.org/10.1177/1092623310379137>; PMID:21037200
42. Hennipman A, Oirschot BAV, Smits J, Rijkssen G and Staal GE, Glycolytic enzyme activities in breast cancer metastases. *Tumour Biol*. 1988;9(5):241-8. <https://doi.org/10.1159/000217568>
43. Wang B, Hsu SH, Frankel W, *et al.*, Stat3-mediated activation of microRNA-23a suppresses gluconeogenesis in hepatocellular carcinoma by down-regulating glucose-6-phosphatase and peroxisome proliferator- activated receptor gamma, coactivator 1 alpha. *Hepatology*. 2012;56(1):186-97. <https://doi.org/10.1002/hep.25632>; PMID:22318941 PMID:PMC3355233
44. Chen L, Yu LJ and Waxman DJ, Potentiation of cytochrome P450/cyclophosphamide-based cancer gene therapy by coexpression of the P450 reductase gene. *Cancer Res*. 1997;57(21):4830-7. PMID:9354446.
45. Chiocca EA and Waxman DJ, Cytochrome P450-based gene therapies for cancer. *Methods Mol Med*. 2004;90:203-22. PMID:14657565
46. Portsmouth D, Hlavaty J and Renner M, Suicide genes for cancer therapy. *Mol Aspects Med*. 2007;28(1):4-41. <https://doi.org/10.1016/j.mam.2006.12.001>; PMID:17306358
47. Chen L and Waxman DJ, Cytochrome P450 gene-directed enzyme prodrug therapy (GDEPT) for cancer. *Curr Pharm Des*. 2002;8(15):1405-16. <https://doi.org/10.2174/1381612023394566>.
48. Kan O, Kingsman S and Naylor S, Cytochrome P450-based cancer gene therapy: current status. *Expert Opin Biol Ther*. 2002;2(8):857-68. <https://doi.org/10.1517/14712598.2.8.857>; PMID:12517265
49. Van Kuilenburg ABP, Dekker HL, Van Den Boget C, Nieboer P, Van Gelder BP and Muijsers AO, Isoforms of human cytochrome c oxidase. Subunit composition and steady-state kinetic properties. *Eur. J. Biochem*. 1991;109(3):615-22. <https://doi.org/10.1111/j.1432-1033.1991.tb16162.x>.
50. Wallace DC, Mitochondria and cancer. *Nat Rev Cancer*. 2012; 12: 685-98. <https://doi.org/10.1038/nrc3365>; PMID:23001348 PMID:PMC4371788

51. Periyasamy K, Sivabalan V, Baskaran K, Kasthuri K and Sakthisekaran D, Cellular metabolic energy modulation by tangeretin in 7, 12-dimethylbenz (a) anthracene-induced breast cancer. *J Biomed Res.* 2016;30(2):134-41. PMID:28276668 PMCID:PMC4820890.
52. Pool-Zobel B, Veeriah S and Bohmer FD, Modulation of xenobiotic metabolising enzymes by anticarcinogens –focus on glutathione S-transferases and their role as targets of dietary chemoprevention in colorectal carcinogenesis. *Mutat Res.* 2005;591(1):74-92. <https://doi.org/10.1016/j.mrfmmm.2005.04.020>; PMID:16083918.
53. King R.S, Teitel CH, Shaddock JG, Casciano DA and Kadlubar FF, Detoxification of carcinogenic aromatic and heterocyclic amines by enzymatic reduction of the N-hydroxy derivative. *Cancer Lett.* 1999;143(2):167-71. [https://doi.org/10.1016/S0304-3835\(99\)00119-6](https://doi.org/10.1016/S0304-3835(99)00119-6).
54. Sato K, Glutathione s-transferase as markers of preneoplasia and neoplasia. *Adv Cancer Res.* 1989;52:205-55. [https://doi.org/10.1016/S0065-230X\(08\)60214-6](https://doi.org/10.1016/S0065-230X(08)60214-6).
55. Russo J, Hu Y, Yang X and Russo IX, Developmental, Cellular, and Molecular Basic of Human Breast Cancer. *J. Of National Cancer Institut Monographs.* 2000;27:17-37. <https://doi.org/10.1093/oxfordjournals.jncimonographs.a024241>.
56. Daniet AO, Folahan OA, Ayele G, Adrian A, Ernest BI, Broderick E, *et al.*, Biological Activity and Mass Spectrometric Analysis of Vernonia amygdalina Fractions. *J Biosci Technol.* 2011;2(3)S:287-304.
57. Inoue Y, Hada TA, Shiraishi K., Hirore H, Hamashima and Kobayashi S, Biphasic effects of Geranylgeraniol, Terpenone and Phytol on the growth of *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* 2005;49(5):1770-4. <https://doi.org/10.1128/AAC.49.5.1770-1774.2005>; PMID:15855494 PMCID: PMC1087616.
58. Sermakkani M and Thangapandian V, GC-MS analysis of Cassia italica leaf methanol extract. *Asian J Pharm Clin Res.* 2012;5(2):90-4.
59. Vivancos M and Moreno JJ,  $\beta$ -Sitosterol modulates antioxidant enzyme response in RAW 264 7 macrophages. *Free Radic Biol Med.* 2005;39(1):91-7. <https://doi.org/10.1016/j.freeradbiomed.2005.02.025>; PMID:15925281
60. Awad AB, Toczek J and Fink CS, Phytosterols decrease prostaglandin release in cultured P388D1/MAB macrophages. *Prostaglandins Leukotrienes Essent. Fatty Acids.* 2004;70(6):511-20. <https://doi.org/10.1016/j.plefa.2003.11.005>; PMID:15120714.
61. Awad AB, Chinnam M, Fink CS and Bradford PG,  $\beta$  -Sitosterol activates Fas signaling in human breast cancer cells. *Phytomedicine.* 2007;14(11):747-54. <https://doi.org/10.1016/j.phymed.2007.01.003>; PMID:17350814.
62. Kumar PP, Kumaravel S and Lalitha C, Screening of antioxidant activity, total phenolics and GC-MS study of *Vitex negundo*. *African J Biochem Res.* 2010;4(7):191-5.

## SUMMARY

- The whole aerial parts of *Pothos scandens* were not evaluated for its anticancer properties though it was used by traditional healers to treat cancer.
- The present study investigated the therapeutic effect of 50% hydro-ethanolic extract of aerial parts of *Pothos scandens* (HEEPS) on key carbohydrate metabolizing enzymes (glycolytic [Hexokinase, Phosphoglucose isomerase], gluconeogenic [Glucose-6-phosphatase, Fructose-1,6-bis phosphatase]) and TCA cycle [Succinate dehydrogenase, Malate dehydrogenase] and xenobiotic marker (Cytochrome P450 reductase, Cytochrome C oxidase and Glutathione -S- transferase) enzymes in DMBA induced experimental mammary carcinoma.
- HEEPS showed significant anticancer activity which was revealed by its ability to restore the enzymes levels which were altered in cancer induced groups to near normal in HEEPS treated groups. The phytochemical analysis of HEEPS also showed the presence of compounds with proved anticancer activity.

## About Authors



**Mr. Jethinlalkhosh J P:** Is working as Quality Control Assistant in Food Corporation of India, Palakkad, Kerala, India. Prior to that, he worked as an Assistant Professor, Department of Biotechnology and Applied Microbiology, St. Thomas College, Pala, Kottayam, Kerala, India. Currently he is pursuing his Ph. D from Department of Biochemistry, Research and Development Centre, Bharathiar University, Coimbatore, Tamil Nadu, India, under the guidance of DR. D Victor Arokia Doss. He is having research experience in the field of Phytochemistry, Pharmacology, Clinical Biochemistry and Toxicology, Molecular Biology and Cancer Biology and is having publications in many international peer reviewed journals.



**Mrs. P Praveena:** Is presently working as Assistant Professor, Department of Biotechnology, Sahrdaya college of Engineering and Technology, Thrissur, Kerala. Currently she is pursuing her Ph. D from Department of Biochemistry, Research and Development Centre, Bharathiar University, Coimbatore, Tamil Nadu, India, under the guidance of DR. D Victor Arokia Doss. Her major research area and publications are in the field of Biochemistry, Clinical toxicology and Phytochemistry.



**Dr. Victor Arokia Doss:** Is working as an Associate Professor in the Department of Biochemistry, P S G Arts and Science College, Coimbatore, Tamil Nadu, India since 1994. He has research experience in the field of Clinical Biochemistry and Toxicology, Molecular Biology, Cancer Biology, Immunology, Protein Chemistry and Bioinformatics. Over the years he has authored and co-authored more than 80 research articles in international peer reviewed journals.

**Cite this article:** JPrasannakumari JJ, Padmam P, Doss VA. Therapeutic Effect of Hydro-Ethanolic Extract of *Pothos scandens* L on key Carbohydrate Metabolizing Enzymes and Xenobiotic Marker Enzymes in DMBA Induced Experimental Mammary Carcinoma. *Indian J of Pharmaceutical Education and Research.* 2017;51(3):427-35.