

# Headspace SPME-GC-MS Analysis and *in silico* Molecular Docking Studies of Phytochemical Compounds Present in *Houttuynia cordata* Thunb

Thokchom Biona<sup>1</sup>, Nameirakpam Bunindro Singh<sup>1</sup>, Ojit Singh Keithellakpam<sup>1</sup>, Sanjai Srinivasan<sup>2</sup>, Shankarrao Patil<sup>2</sup>, Pulok Kumar Mukherjee<sup>1</sup>, Nanaocha Sharma<sup>1,\*</sup>

<sup>1</sup>Institute of Bioresources and Sustainable Development, Takyelpat, Imphal, Manipur, INDIA

<sup>2</sup>Dhriti Global Education Network, Bangalore, Karnataka, INDIA.

## ABSTRACT

**Aim:** The study aimed to identify the phytochemical constituents present in the leaves, stems, and roots of *H. cordata* and perform *in silico* molecular docking studies of selected common compounds present in all three parts. **Materials and Methods:** The phytochemical components were investigated using the headspace solid-phase micro-extraction followed by gas chromatography-mass spectrometry and molecular docking was performed using Autodock Vina v.1.2.0. **Results:**  $\beta$ -pinene was found to be the major compound present in stem (73.89%), leaves (66.46%) and roots (42.88%). Four category targets were used for *in silico* molecular docking of 14 common compounds found in leaves, roots, and stems. The present study showed that the compounds caryophyllene and dihydro-cis-alpha-copaene-8-ol had antibacterial, antioxidant, anti-cancer, and anti-inflammatory activities. **Conclusion:** This work demonstrated the great utility of headspace SPME-GC-MS for the investigation of aromatic chemicals in a variety of edible and medicinal spices. With *in silico* molecular docking, we may look into the potential pharmacological activity of various volatile organic compounds present in *H. cordata*.

**Keywords:** *Houttuynia cordata*, Headspace SPME-GC-MS, *in silico* molecular docking,  $\beta$ -pinene, Caryophyllene.

## Correspondence:

**Dr. Nanaocha Sharma**

Institute of Bioresources and Sustainable Development, Takyelpat, Imphal-795001 Manipur, INDIA.

Email: sharma.nanaocha@gmail.com

**Received:** x-x-x;

**Revised:** x-x-x;

**Accepted:** x-x-x.

## INTRODUCTION

Indigenous foods not only have a distinct flavor and taste of their own, but they also offer dietary variety and ensure the security of the household's food supply. The diets of the northeastern peoples of India are primarily composed of natural and traditional foods. *H. cordata*, locally referred to as "Toningkhok," is one such native food plant of Manipur. *H. cordata* is a perennial, aromatic herb belonging to the family Saururaceae, a fragrant medicinal plant with spreading rootstock. The Manipuri people use the herb both raw and cooked, and the local healer (Maiba) uses it to treat anaemia, gastritis, and tuberculosis as well as dysentery, muscular sprains, and stomach ulcers. Alkaloids, essential oils, flavonoids, and other chemical components with distinctive therapeutic effects are among the chemical ingredients found in *H. cordata*. In Manipur's plain and hill districts, the plant grows wild and is harvested for market sale. In Meghalaya, it is used

in salads or cooked with other vegetables.<sup>1</sup> Leaf juices are used to cure cholera, dysentery, treat blood deficiencies, and blood purification.<sup>2</sup> Naga tribe of Kiphire District, Nagaland uses the whole plant to cure stomach ache, cholera, and dysentery and as diuretic. It is also applied to skin diseases.<sup>3</sup> Apatani Tribe in Arunachal Pradesh used the shoot of *H. cordata* for freshness, good sleep, and heart disorders.<sup>4</sup> In the Senapati district of Manipur, roots and leaves are used to cure measles, gonorrhoea, and skin troubles.<sup>5</sup> In Southeast and East Asian countries, it is frequently used as an herbal anti-inflammatory, antibacterial, antiviral, and anti-cancer medicine. *H. cordata* has the potential for both antiviral activity and antiviral consequences, which is relevant given the recent COVID-19 pandemic reported by Lau *et al.* (2008) during the SARS outbreak of 2002-2003.<sup>6</sup> Recently, several studies have also produced scientific evidence to support and reveal its anti-inflammatory, anti-allergic, virucidal, anti-oxidative and anti-cancer activities.<sup>7-12</sup> Due to its quick and ease of use, Headspace SPME is useful in the identification of plant volatile profiles. In addition, it acts as a tool to distinguish the phytochemicals present in different parts of the plant of the same species or different species.



DOI: 10.5530/ijper.58.3.95

### Copyright Information :

Copyright Author (s) 2024 Distributed under Creative Commons CC-BY 4.0

Publishing Partner : EManuscript Tech. [www.emanuscript.in]

## MATERIALS AND METHODS

### Sample Collection and Preparation

The fresh portions of *H. cordata* were collected in September 2022 from Takyelpat, Imphal West District, Manipur. The plant was identified and authenticated by Botanical Survey of India, Sikkim Himalayan Regional Centre with voucher number IBSD-SC/EPS/2020/IP/123. The authenticated herbarium was submitted in the IBSD Herbarium Library. The three parts of *H. cordata* viz., fresh leaves, roots, and stems were packed in a 15 mL clear vial which has a screw-top hole cap with silicone septa. All the vials were filled with one-third portions of the sample. The roots and stems were cut into identical lengths of 2 cm.

### Headspace SPME-GC-MS

The SPME fiber was conditioned in the GC-MS and confirmed with no impurities through the blank run. The 15 mL glass vial containing the sample was exposed to the SPME fiber for 20 min. Each sample after the exposure was injected into the inlet injector of the GC-MS instrument for 2 min. Trace 1300 GC fitted with TG-5MS column and mass detector (TSQ DUO) with triple Quadropole was used to study the volatile organic compounds. Helium was used as the carrier gas, with a flow rate of 1 mL/min, and a split ratio of 1:20 was maintained. The initial column temperature was programmed from 40°C for 1 min to 250°C at a rate of 5°C per min, and then to 250°C for 20 min.<sup>13,14</sup> The relative VOC's constituents were expressed by its peak area percentage. Based on the comparison of mass spectra with those of the 2017 National Institute of Standards and Technology (NIST) GC-MS Libraries the volatile compounds were identified.

### 3D structure retrieval of protein targets

The Anti-Microbial (AM), Anti-Oxidant (AO), Anti-Cancer (AC), and Anti-Inflammatory (AI) targets were selected based on their prominence in the literature and their 3D structure was retrieved from Protein Data Bank (<https://www.rcsb.org/>).

### 3D ligand structure retrieval

The structure of 1,7-Octadiene, 2-methyl-6-methylene was downloaded from NIST Chemistry WebBook (<https://webbook.nist.gov/>).<sup>15</sup> All the other ligand structures were downloaded from PUBCHEM database (<https://pubchem.ncbi.nlm.nih.gov/>).<sup>16</sup>

### Protein and Ligand preparation

Auto Dock MGL Tools v.1.5.7 was used for the protein preparation step.<sup>17</sup> The identical chains, the native co-crystallized ligands, and the water molecules were deleted at first, the protein structure was added with polar hydrogen atoms and the Gasteiger charges were introduced. For the preparation of ligand molecules Chimera v.1.1.6 software was utilized, the ligands were protonated and the charges were added.<sup>18</sup> For the file format conversions Openbabel tool was used.<sup>19</sup>

### Binding site prediction

The binding sites of Antimicrobial and Antioxidant targets were comprehensively documented in previous research were used.<sup>20</sup> For Anticancer and Anti-inflammatory targets, the binding site grid coordinates were predicted by using BIOVIA Discovery Studio, at first the co-crystallised ligands were selected and the "Define and Edit Binding Site" option was used.<sup>21</sup>

### Molecular docking and analysis

Autodock Vina v.1.2.0 which uses the Lamarckian algorithm was used for performing molecular docking.<sup>22</sup> The native co-crystallized ligands and the short-listed phytochemical compounds were docked individually using the command line. For saving the docked ligand and protein complexes Pymol software was used.<sup>23</sup> The images of active pocket visualization and 2D amino acid interaction between the ligand-protein complexes were obtained from BIOVIA Discovery Studio.

## RESULTS

### Phytochemical analysis using Headspace SPME-GC-MS

A total of 36, 30, and 25 compounds were identified in the fresh leaves (97.9%), roots (97.55%), and stems (98.30%) of *H. cordata* respectively (Tables 1, 2, and 3). The GCMS Chromatogram of all three parts of *H. cordata* is shown in Figure 1. The major compounds present in the fresh leaves of *H. cordata* were  $\beta$ -pinene (66.46%), Leaf alcohol (6.05%),  $\beta$ -Sabinene (2.36%), Caryophyllene (2.21%), Camphene (2.21%),  $\zeta$ -Terpinene (1.91%), o-Cymene (1.81%), 2-Norpinene, 3,6,6-trimethyl- (1.8%), 1,7-Octadiene 2-methyl-6-methylene (1.4%), Cyclohexene, 1-methyl-4-(1-methylethylidene)-(1.36%). For fresh roots of *H. cordata*, the dominant compounds were  $\beta$ -pinene (42.88%), m-Mentha-6,8-diene, (R)-(+)- (12.88%), p-Mentha-1(7),3-diene (8.04%),  $\alpha$ -Pinene (7.06%), Bicyclo[3.1.0]hexane, 4-methylene-1-(1-methylethyl)-(4.82%), o-Cymene (3.27%),  $\zeta$ -Terpinene (2.81%), Cyclohexene, 1,5,5-trimethyl-3-methylene (2.5%), 2-Thujene (2.45%), p-Mentha-1,4(8)-diene (2.14%), Camphene (1.29%), 1,3-Cyclohexadiene, 1,3,5,5-tetramethyl (1.1%) and Caryophyllene (1.07%). Other major compounds found in stem are Sabinene (3.61%), D-sylvestrene (2.47%),  $\beta$ -Terpinene (2.45%), Caryophyllene (1.92%), Bornyl acetate (1.91%),  $\beta$ -Thujene (0.95%), Camphene (0.94%) and o-Cymene (0.91%). The volatile organic compounds present in the leaves, roots, and stems were predominantly monoterpenoids. Monoterpenes have antimicrobial, anti-inflammatory, antioxidant, antipruritic, hypotensive, and analgesic pharmacological properties. Terpenoids are significant for more reasons than just their aroma; they are also linked to anti-insect action and positive aspects of human health. For *in silico* studies, 14 active compounds prevalent in all the three samples i.e. leaves, roots and stem were chosen and further used for *in silico* studies.

## Molecular docking analysis

The Binding site grid coordinates used for molecular docking is listed in Table 4. The molecular docking of 14 phytochemical compounds with various targets was performed. The binding energies of all the compounds are represented as heat map as shown in Figure 2. The compounds Caryophyllene and dihydro-cis-alpha-copaene-8-ol were highlighted exclusively in this study as both of them had binding affinity to a higher number of targets compared to other compounds. Table 5 contains the binding energies and amino acid interactions of phytochemical compounds Caryophyllene and dihydro-cis-alpha-copaene-8-ol against various targets.


The compounds caryophyllene and dihydro-cis-alpha-copaene-8-ol showed high binding affinity of -7.7 and -7.8 kcal/mol against the antimicrobial target dihydrofolate reductase of *S. aureus* (PDB-ID: 3SRW). The enzyme Dihydrofolate Reductase (DHFR) is responsible for the NADPH-dependent conversion of dihydrofolate to tetrahydrofolate.<sup>24</sup> Tetrahydrofolate is essential for several biosynthetic pathways, including amino acid and nucleic acid metabolism.<sup>25</sup> DHFR inhibitors are effective for treating bacterial, mycobacterial, fungal, and protozoal infections hence we chose DHFR as an anti-microbial target.<sup>25</sup> Caryophyllene interacted with the target DHFR with 9 van der Waals interactions and 9 Pi interactions. dihydro-cis-alpha-copaene-8-ol interacted with 1 hydrogen bond, 4 van der Waals interactions, and 10 pi

interactions. The amino acid interactions of the compounds to the anti-microbial target 3SRW are shown in Figure 3.

Both compounds showed high binding affinities of -7.4 and -7.1 kcal/mol against the antioxidant target cytochrome P450 CYP2C9 (PDBID: 1OG5). Cytochrome P450 enzymes produce reactive oxygen species which provide oxidative stress, inhibiting this enzyme produces an antioxidant effect.<sup>26,27</sup> Caryophyllene interacted with the target CYP2C9 with 4 van der Waals interactions and 10 Pi interactions. dihydro-cis-alpha-copaene-8-ol interacted with 1 hydrogen bond, 11 van der Waals interactions, and 10 pi interactions. The amino acid interactions of the compounds to the antioxidant target 1OG5 are shown in Figure 4.

The compounds showed higher binding affinities of -8.5 and -8 kcal/mol against the anticancer target Human Estrogen Receptor Alpha (PDBID: 3ERT) than other anticancer targets. The Estrogen Receptor alpha (ER $\alpha$ ) is known to play an important role in cell proliferation in Breast cancer hence we used ER $\alpha$  as a potential anti-cancer target.<sup>28,29</sup> Caryophyllene interacted with the target 3ERT with 7 van der Waals interactions and 8 Pi interactions. dihydro-cis-alpha-copaene-8-ol interacted with 6 van der Waals interactions and 9 pi interactions. The amino acid interactions of the compounds to the anticancer target 3ERT are shown in Figure 5 (A).

Though the compounds were having binding affinity to other anti-inflammatory targets, TNF-alpha was particularly chosen due to its extensive usage in the literature as an anti-inflammatory



**Figure 1:** (A) Chromatogram of Headspace SPME-GC-MS *H. cordata* leaves, (B) Chromatogram of Headspace SPME-GC-MS *H. cordata* roots. (C) Chromatogram of Headspace SPME-GC-MS *H. cordata* stem.

**Table 1: Major volatile organic compounds present in *Houttuynia cordata* leaves.**

| Sl. No. | RT <sup>a</sup> | Compound   | RSI <sup>b</sup> | RA <sup>c</sup> % | Mol. Wt |
|---------|-----------------|--|------------------|-------------------|---------|
| 1       | 9.38            | β-pinene   | 894              | 66.46             | 136     |
| 2       | 5.73            | Leaf alcohol   | 911              | 6.05              | 100     |
| 3       | 8.76            | β-Sabinene   | 944              | 2.36              | 136     |
| 4       | 21.27           | Caryophyllene  | 941              | 2.21              | 204     |
| 5       | 10.36           | Camphene   | 872              | 2.21              | 136     |
| 6       | 11.25           | ç-Terpinene  | 905              | 1.91              | 136     |
| 7       | 10.25           | o-Cymene   | 920              | 1.81              | 134     |
| 8       | 10.64           | 2-Norpinene, 3,6,6-trimethyl-  | 897              | 1.80              | 136     |
| 9       | 8.93            | 1,7-Octadiene, 2-methyl-6-methylene  | 885              | 1.40              | 136     |
| 10      | 10.01           | p-Menth-4(8)-ene   | 926              | 1.36              | 136     |
| 11      | 12.12           | (+)-4-Carene   | 885              | 1.00              | 136     |
| 12      | 7.46            | β-Thujene  | 931              | 0.94              | 136     |
| 13      | 15.48           | Decanal  | 878              | 0.75              | 156     |
| 14      | 10.94           | á-Ocimene  | 908              | 0.72              | 136     |
| 15      | 13.35           | 2,6-Dimethyl-1,3,5,7-octatetraene, E,E   | 909              | 0.67              | 134     |
| 16      | 13.68           | Alloocimene  | 947              | 0.65              | 136     |
| 17      | 17.36           | 1-Decanol  | 928              | 0.63              | 158     |
| 18      | 22.94           | [1,4]Dioxino[2,3-b]-1,4-dioxin, hexahydro-2,2,3,3-tetramethyl                                  | 826              | 0.52              | 202     |
| 19      | 7.64            | 3-Carene   | 922              | 0.49              | 136     |
| 20      | 17.92           | gamma-Terpinene diepoxide  | 928              | 0.47              | 168     |
| 21      | 8.84            | 3-p-Menthene   | 931              | 0.41              | 136     |
| 22      | 6.04            | Cyclopropane, propyl-  | 879              | 0.36              | 84      |
| 23      | 9.67            | α-Thujene  | 913              | 0.34              | 136     |
| 24      | 23.15           | 1H-Cyclopropa[a]naphthalene, decahydro-1,1,3a-trimethyl-7-methylene-, [1aS-(1aà,3aà,7aá,7bà)]- | 892              | 0.32              | 204     |
| 25      | 14.74           | 3-Cyclohexen-1-ol, 4-methyl-1-(1-methylethyl)-, (R)-   | 895              | 0.30              | 154     |
| 26      | 5.62            | 2-Hexenal  | 909              | 0.28              | 98      |
| 27      | 9.80            | (Z),(Z)-2,4-Hexadiene  | 861              | 0.26              | 82      |
| 28      | 10.09           | Cyclopropane, 1-ethenyl-2-hexenyl-, [1à,2á(E)]-(ñ)-  | 849              | 0.25              | 150     |
| 29      | 22.12           | Cyclohexene, 4-[(1E)-1,5-dimethyl-1,4-hexadien-1-yl]-1-methyl                                  | 893              | 0.15              | 204     |
| 30      | 7.10            | Furan, 2-ethyl-  | 882              | 0.15              | 96      |
| 31      | 26.47           | 2-Hydroxy-1-(1'-pyrrolidiyl)-1-buten-3-one   | 799              | 0.14              | 155     |
| 32      | 8.48            | 3,5-Dimethylamphetamine  | 889              | 0.13              | 163     |
| 33      | 13.10           | 2,6-Dimethyl-1,3,5,7-octatetraene, E,E   | 902              | 0.11              | 134     |
| 34      | 14.04           | 2-Hydroxymandelic acid, ethyl ester, di-TMS  | 880              | 0.10              | 340     |
| 35      | 25.49           | 2,2,4-Trimethyl-1,3-pentanediol diisobutyrate  | 862              | 0.10              | 286     |
| 36      | 21.10           | Dihydro-cis-à-copaene-8-ol   | 905              | 0.09              | 222     |
|         |                 | Total area %   |                  | 97.9              |         |

RT<sup>a</sup>: Retention time; RSI<sup>b</sup>: Reversed search index on TG-5MS capillary column, RA<sup>c</sup> %: Relative area (peak area relative to the total peak area).

**Table 2: Major volatile organic compounds present in *Houttuynia cordata* roots.**

| Sl. No. | RT <sup>a</sup> | Compound  | RSI <sup>b</sup> | RA <sup>c</sup> % | Mol.wt |
|---------|-----------------|---|------------------|-------------------|--------|
| 1       | 9.36            | β-pinene  | 884              | 42.88             | 136    |
| 2       | 10.39           | D-sylvestrene   | 880              | 12.88             | 136    |
| 3       | 8.87            | p-Mentha-1(7),3-diene   | 925              | 8.04              | 136    |
| 4       | 7.66            | alpha-Pinene  | 929              | 7.06              | 136    |
| 5       | 8.78            | Sabinene  | 945              | 4.82              | 136    |
| 6       | 10.27           | o-Cymene  | 919              | 3.27              | 134    |
| 7       | 11.26           | ζ-Terpinene   | 900              | 2.81              | 136    |
| 8       | 10.02           | Cyclohexene, 1,5,5-trimethyl-3-methylene  | 894              | 2.5               | 136    |
| 9       | 7.48            | 2-Thujene   | 921              | 2.45              | 136    |
| 10      | 12.13           | p-Mentha-1,4(8)-diene   | 937              | 2.14              | 136    |
| 11      | 8.06            | Camphene  | 944              | 1.29              | 136    |
| 12      | 13.67           | 1,3-Cyclohexadiene, 1,3,5,5-tetramethyl   | 921              | 1.1               | 136    |
| 13      | 21.27           | Caryophyllene   | 942              | 1.07              | 204    |
| 14      | 13.31           | Alloocimene   | 931              | 0.86              | 136    |
| 15      | 9.67            | 3-Thujene   | 900              | 0.79              | 136    |
| 16      | 8.94            | 1,7-Octadiene, 2-methyl-6-methylene   | 874              | 0.75              | 136    |
| 17      | 10.65           | 2-Norpinene, 3,6,6-trimethyl  | 900              | 0.67              | 136    |
| 18      | 11.26           | á-Ocimene   | 935              | 0.64              | 136    |
| 19      | 17.78           | Bornyl acetate  | 932              | 0.51              | 196    |
| 20      | 14.75           | 3-Cyclohexen-1-ol, 4-methyl-1-(1-methylethyl)-, (R)-                                | 897              | 0.32              | 154    |
| 21      | 17.91           | 4,8-Dioxatricyclo[5.1.0.0(3,5)] octane, 1-methyl-5-(1-methylethyl)-, (1à,3á,5á,7à)- | 906              | 0.27              | 168    |
| 22      | 9.59            | 2,6-Dimethyl-2-trans-6-octadiene  | 844              | 0.15              | 138    |
| 23      | 11.58           | 2-Cyclohexen-1-ol, 4-ethyl-1,4-dimethyl   | 854              | 0.15              | 154    |
| 24      | 14.44           | Borneol   | 917              | 0.13              | 154    |
| 25      | 9.84            | 1-Propanone, 1-(1-cyclohexen-1-yl)-   | 905              | 0.11              | 138    |
|         |                 | Total Area %  |                  | 97.55             |        |

RT<sup>a</sup>: Retention time; RSI<sup>b</sup>: Reversed search index on TG-5MS capillary column, RA<sup>c</sup> %: Relative area (peak area relative to the total peak area).

| List of shortlisted compounds                        | Anti-microbial targets |      |      |      |      | Anti-oxidant targets |      |      |      | Anti-cancer targets |      |      |       |      | Anti-inflammatory targets |      |       |      |      |
|--|------------------------|------|------|------|------|----------------------|------|------|------|---------------------|------|------|-------|------|---------------------------|------|-------|------|------|
|  | 1JZQ                   | 1KZN | 2ZDQ | 3SRW | 3TIZ | 1N8Q                 | 1OG5 | 2CDU | 3NRZ | 3ERT                | 1D18 | 1M17 | 3OG7  | 6JOK | 4O1Z                      | 5KIT | 3w5c  | 5KX7 | 2AZ5 |
| Co-crystallised ligands                              | -8.9                   | -8.5 | -7   | -9.7 | -7.7 | -5.6                 | -9.4 | -8   | -6.1 | -10                 | -8.8 | -6.3 | -10.1 | -7.4 | -8.2                      | -8.8 | -11.3 | -7   | -9.2 |
| 1,7-Octadiene, 2-methyl-6-methylene                  | -5.3                   | -5.1 | -5.1 | -2.3 | -5.1 | -5.6                 | -5.3 | -4.4 | -6   | -5                  | -5.5 | -4.2 | -5.3  | -5.7 | -5.5                      | -5.7 | -6.2  | -5.8 | -4.8 |
| á-Ocimene  | -5.2                   | -5.2 | -5.3 | -5   | -5.2 | -5.3                 | -5.3 | -4.5 | -6.2 | -5.3                | -6   | -4.6 | -5.6  | -5.9 | -5.8                      | -5.9 | -6.5  | -6   | -5   |
| Alloocimene  | -5.5                   | -5.5 | -5.3 | -5.3 | -5.6 | -5.2                 | -5.7 | -4.7 | -6.3 | -5.4                | -6.3 | -4.9 | -5.8  | -5.8 | -5.8                      | -5.9 | -6.8  | -5.8 | -5.2 |
| 3-Cyclohexen-1-ol, 4-methyl-1-(1-methylethyl)-, (R)- | -5.9                   | -6.1 | -6   | -5.9 | -5.4 | -5.8                 | -5.9 | -4.4 | -4.4 | -6                  | -6.5 | -5.2 | -6.1  | -5.7 | -6                        | -5.8 | -6.4  | -6.4 | -5.6 |
| β-pinene   | -5.3                   | -4.6 | -4.9 | -5.5 | -4.9 | -4.9                 | -5.6 | -3.4 | -1.1 | -6.6                | -5.9 | -5.2 | -5.7  | -5.6 | -6.1                      | -5.4 | -6    | -5.8 | -5.6 |
| Caryophyllene  | -6.5                   | -5.6 | -5.9 | -7.7 | -6   | -2.1                 | -7.4 | -4.3 | 2    | -8.5                | -7.7 | -6.9 | -7.4  | -5.9 | -7.9                      | -5   | -7.8  | -6.6 | -7   |
| Camphene   | -5.2                   | -4.7 | -4.7 | -5.5 | -4.7 | -3.7                 | -5.6 | -3.3 | 0.3  | -6.4                | -5.9 | -5.1 | -5.8  | -5.4 | -6.8                      | -5.4 | -6.5  | -5.6 | -5.5 |
| ζ-Terpinene  | -5.5                   | -5.8 | -5.5 | -5.6 | -5.7 | -5.6                 | -6.1 | -3.9 | -6.8 | -5.8                | -6.7 | -5.3 | -6.2  | -6.3 | -6.2                      | -6.5 | -7.1  | -6.5 | -5.5 |
| o-Cymene   | -5.1                   | -5.5 | -5.3 | -5.5 | -5.3 | -5.9                 | -5.8 | -3.4 | -7   | -5.8                | -6.3 | -5.3 | -6.2  | -6.4 | -6.4                      | -6.5 | -7    | -6.1 | -5.4 |
| β-Thujene  | -5                     | -5   | -5.1 | -5.4 | -5   | -5.1                 | -5.4 | -5   | -3.5 | -5.8                | -6.1 | -5.1 | -6.2  | -6.5 | -6                        | -6   | -6.3  | -6.1 | -5.5 |
| 2,6-Dimethyl-1,3,5,7-octatetraene, E,E               | -5.5                   | -5.5 | -5.3 | -5.2 | -5.7 | -5.5                 | -5.8 | -4.6 | -6.3 | -5.5                | -6.4 | -4.9 | -5.8  | -6.1 | -5.7                      | -6   | -6.8  | -5.7 | -5.3 |
| Dihydro-cis-á-copaene-8-ol                           | -6.7                   | -6.2 | -4.6 | -7.8 | -6.8 | -0.1                 | -7.1 | -4.2 | 2    | -8                  | -7.7 | -6.8 | -7.6  | -5.2 | -7.2                      | -2.3 | -7.5  | -6.9 | -7.8 |
| D-sylvestrene  | -5.2                   | -5.9 | -5.4 | -5.7 | -5.6 | -4.8                 | -6   | -4.2 | -6.7 | -5.9                | -6.5 | -5.3 | -6.2  | -6.5 | -6.3                      | -6.7 | -7.3  | -6   | -5.7 |
| Bornyl acetate                                       | -5.8                   | -5.2 | -5.5 | -6.5 | -5.6 | -1                   | -6.1 | -3.9 | 0.8  | -6.7                | -6.2 | -5.3 | -5.5  | -4.9 | -6.8                      | -3.5 | -5.8  | -4.7 | -6.5 |

**Figure 2:** Heat-map representation of binding energies (in kcal/mol) of Phytochemical compounds against various targets.



**Table 3: Major volatile organic compounds present in *Houttuynia cordata* stem.**

| Sl. No. | RT <sup>a</sup> | Compound name  | RSI <sup>b</sup> | RA <sup>c</sup> % | Mol. Wt |
|---------|-----------------|--|------------------|-------------------|---------|
| 1       | 9.17            | β-pinene   | 905              | 73.89             | 136     |
| 2       | 8.64            | β-Terpinene  | 930              | 3.61              | 136     |
| 3       | 7.52            | alpha-pinene   | 933              | 3.48              | 136     |
| 4       | 10.23           | D-sylvestrene  | 882              | 2.47              | 136     |
| 5       | 8.72            | β-Terpinene  | 939              | 2.45              | 136     |
| 6       | 21.12           | Caryophyllene  | 919              | 1.92              | 204     |
| 7       | 17.60           | Bornyl acetate   | 912              | 1.91              | 196     |
| 8       | 7.35            | β-Thujene  | 915              | 0.95              | 136     |
| 9       | 7.93            | Camphene   | 947              | 0.94              | 136     |
| 10      | 10.12           | o-Cymene   | 931              | 0.91              | 134     |
| 11      | 11.12           | ç-Terpinene  | 892              | 0.88              | 136     |
| 12      | 17.04           | 6-(3-Methyl-3-cyclohexenyl)- 2-methyl-2,6-heptadienol          | 877              | 0.88              | 220     |
| 13      | 9.88            | α-Terpinolene  | 897              | 0.48              | 136     |
| 14      | 11.99           | α-Terpinolene  | 889              | 0.39              | 136     |
| 15      | 17.87           | Heptane, 3,3-dimethyl-   | 904              | 0.32              | 128     |
| 16      | 9.96            | Diazoadamantane  | 890              | 0.29              | 162     |
| 17      | 22.99           | 10-epi-ç-Eudesmol  | 885              | 0.29              | 222     |
| 18      | 16.70           | Phenylacetaldehyde N-methyl-N-formylhydrazone                  | 829              | 0.28              | 176     |
| 19      | 22.22           | Cyclohexene, 1,5,5-trimethyl-6-(2-propenylidene)-              | 936              | 0.26              | 162     |
| 20      | 13.54           | 2,4,6-Octatriene, 2,6-dimethyl-                                | 867              | 0.25              | 136     |
| 21      | 13.20           | 2,6-Dimethyl-1,3,5,7-octatetraene, E,E                         | 825              | 0.23              | 134     |
| 22      | 10.51           | 4-Carene, (1S,3R,6R)-(-)-                                      | 884              | 0.20              | 136     |
| 23      | 22.81           | Dihydro-cis-α-copaene-8-ol                                     | 938              | 0.17              | 222     |
| 24      | 21.97           | Camphene   | 850              | 0.16              | 136     |
| 25      | 10.81           | Tricyclo[3.2.1.0(2,4)]octane, 8-methylene-, (1à,2à,4à,5à)-     | 925              | 0.14              | 120     |
| 26      | 14.58           | 2-Cyclopenten-1-one, 2,3,5-trimethyl-4-methylene               | 889              | 0.13              | 136     |
| 27      | 9.52            | β-Thujene  | 878              | 0.12              | 136     |
| 28      | 12.09           | Tricyclo[3.2.1.0(2,,4)]oct-6-ene, 8-methylene-, (1à,2à,4à,5à)- | 891              | 0.10              | 118     |
| 29      | 14.27           | 2,8-Bornanediol  | 925              | 0.10              | 170     |
| 30      | 17.75           | 2-Nonanone, 3-(hydroxymethyl)-                                 | 888              | 0.10              | 172     |
|         |                 | Total Area %   |                  | 98.30             |         |

RT<sup>a</sup>: Retention time; RSI<sup>b</sup>: Reversed search index on TG-5MS capillary column, RA<sup>c</sup> %: Relative area (peak area relative to the total peak area).


target, making it an ideal selection for this study. The compounds showed high binding affinities of -7 and -7.8 kcal/mol against the anti-inflammatory Target Tumor Necrosis factor (TNF-alpha) (PDBID: 2AZ5). TNF-alpha is a pro-inflammatory cytokine that plays a major role in the pathogenesis of several inflammatory diseases. We chose TNF-alpha as a potential anti-inflammatory target because TNF-alpha inhibitors are used in the treatment

of Rheumatoid arthritis, Inflammatory Bowel Disease, Psoriasis, and other inflammatory diseases.<sup>30-32</sup> Caryophyllene interacted with the target TNF-alpha with 10 van der Waals interactions and 7 Pi interactions. dihydro-cis-alpha-copaene-8-ol interacted with the target 3SRW with 10 van der Waals interactions and 5 pi interactions. The amino acid interactions of the compounds to the anti-inflammatory target 2AZ5 are shown in Figure 5 (B).

**Table 4: Binding site grid coordinates of targets used for molecular docking.**


| Target Type             | PDB ID | Target Name                                    | Grid centre coordinates |
|-------------------------|--------|--|-------------------------|
| Anti-microbial targets. | 1JZQ   | Isoleucyl-tRNA synthetase.                     | -26.73582776x           |
|                         |        |  | 6.926711078y            |
|                         |        |  | -27.82592825z           |
|                         | 1KZN   | DNA GYRASE                                     | 19.46390268x            |
|                         |        |  | 31.38737131y            |
|                         |        |  | 36.35869076z            |
|                         | 2ZDQ   | D-alanine--D-alanine ligase.                   | 48.35624583x            |
|                         |        |  | 18.85051502y            |
|                         |        |  | -1.467031607z           |
|                         | 3SRW   | Dihydrofolate reductase.                       | -5.437161837x           |
|                         |        |  | -31.03416816y           |
|                         |        |  | 5.382902144z            |
|                         | 3TTZ   | DNA gyrase subunit B.                          | 15.59966623x            |
|                         |        |  | -18.15613991y           |
|                         |        |  | 7.092968912z            |
| Anti-oxidant targets.   | 1N8Q   | Lipoxygenase                                   | 22.36296039x            |
|                         |        |  | 1.272871124y            |
|                         |        |  | 20.2650223z             |
|                         | 1OG5   | CYP2C9   | -19.82366963x           |
|                         |        |  | 86.69793369y            |
|                         |        |  | 38.27579945z            |
|                         | 2CDU   | NADPH-oxidase                                  | 18.99749909x            |
|                         |        |  | -5.670402997y           |
|                         |        |  | -1.718618562z           |
|                         | 3NRZ   | Xanthine oxidase                               | 37.47367438x            |
|                         |        |  | 19.30785549y            |
|                         |        |  | 18.15215059             |
| Anticancer targets      | 3ERT   | Estrogen receptor alpha.                       | 31.574552x              |
|                         |        |  | -1.590379y              |
|                         |        |  | 25.599483z              |
|                         | 1DI8   | Cyclin-dependent kinase 2.                     | -8.773273x              |
|                         |        |  | 50.057273y              |
|                         |        |  | 12.795045z              |
|                         | 1M17   | Epidermal growth factor receptor.              | 22.01369x               |
|                         |        |  | 0.252828y               |
|                         |        |  | 52.794034z              |
|                         | 3OG7   | AKAP9-BRAF fusion protein.                     | 1.868515x               |
|                         |        |  | -2.637667y              |
|                         |        |  | -19.917727z             |
|                         | 6JOK   | Platelet-derived growth factor receptor alpha. | 17.588379x              |
|                         |        |  | 132.559517y             |
|                         |        |  | -6.030276z              |

| Target Type                | PDB ID    | Target Name                                 | Grid centre coordinates |
|----------------------------|-----------|---|-------------------------|
| Anti-inflammatory targets. | 4O1Z      | Cyclooxygenase-1                            | 252.107414x             |
|                            |           |   | 106.979717y             |
|                            |           |   | 4.707935z               |
|                            | 5KIT      | Nicotinamide phosphoribosyltransferase.     | 165.421241x             |
|                            |           |   | 185.734552y             |
|                            |           |   | 192.381517z             |
|                            | 3w5e      | Phosphodiesterase 4B                        | 24.701364x              |
|                            |           |   | 18.137182y              |
|                            |           |   | -18.425333z             |
|                            | 5KX7      | Interleukin-1 receptor-associated kinase 4. | 33.833333x              |
| 40.25825y                  |           |   |                         |
| 59.695417z                 |           |   |                         |
| 2AZ5                       | TNF-alpha | -19.40960x                                  |                         |
|                            |           | 74.650750y                                  |                         |
|                            |           | 33.849550z                                  |                         |




**Figure 3:** Docking results depiction of the key amino acid interactions in the binding site of anti-microbial target DHFR (PDBID: 3SRW) with Caryophyllene and dihydro-cis-alpha-copaene-8-ol.





**Figure 4:** Docking results depiction of the key amino acid interactions in the binding site of anti-oxidant target CYP2C9 (PDBID: 1OG5) with Caryophyllene and dihydro-cis-alpha-copaene-8-ol.



**Figure 5:** (A) Docking results depiction of the key amino acid interactions in the binding site of anti-cancer target ER $\alpha$  (PDBID: 3ERT) with Caryophyllene and dihydro-cis-alpha-copaene-8-ol. (B) Docking results depiction of the key amino acid interactions in the binding site of anti-inflammatory target TNF-alpha (PDBID: 2A25) with Caryophyllene and dihydro-cis-alpha-copaene-8-ol.

**Table 5: Binding energies and amino acid interactions of phytochemical compounds Caryophyllene and dihydro-cis-alpha-copaene-8-ol against various targets.**

| Targets           | PDB-ID | Compounds                          | Binding energy (kcal/mol) | H-bonds                      | Vander Valls interactions   | Pi interactions   |
|-------------------|--------|------------------------------------|---------------------------|------------------------------|---|---|
| Antimicrobial     | 3SRW   | Native ligand - CID 53346511       | -9.7                      | LEU-6, ALA-8, ASP-48, PHE-93 | VAL-7, GLN-20, HIS-31, THR-47, SER-50, GLY-94, PHE-99, THR-112.   | LEU-6, ALA-8, LEU-21(2), LEU-29(2), VAL-32(2), ILE-51(2), LEU-55, PHE-93(3)         |
|                   |        | Caryophyllene                      | -7.7                      | -                            | LEU-6, GLY-16, GLN-20, TRP-23, VAL-32, THR-47, GLY-94, GLY-95, PHE-99.  | VAL-7, ALA-8, ILE-15, LEU-21(5), PHE-93   |
|                   |        | dihydro-cis-alpha-copaene-8-ol     | -7.8                      | PHE-93                       | VAL-7, ASP-28, GLY-94, GLY-95.  | LEU-6, ALA-8, ILE-15, LEU-21, VAL-32(2), PHE-93(3), PHE-99.                         |
| Antioxidant       | 1OG5   | Native ligand - CID 54688261       | -9.4                      | ASN-217                      | GLY-98, LEU-102, VAL-113, ILE-213, GLN-214, THR-364, SER-365, LEU-388.  | ARG-97, ILE-99, PHE-100, ALA-103(2), PHE-114(2), LEU-366, PRO-367(2), PHE-476.      |
|                   |        | Caryophyllene                      | -7.4                      | -                            | ILE-99, LEU-208, ASN-217, PHE-476.  | PHE-100, LEU-102, ALA-103(3), PHE-114, LEU-366(2), PRO-367(2).                      |
|                   |        | dihydro-cis-alpha-copaene-8-ol     | -7.1                      | ASN-217                      | PHE-100, LEU-102, LEU-208, GLN-214, THR-364, SER-365, GLY-475.  | ALA-103, ILE-213, LEU-366(2), PRO-367(5), PHE-476(2).                               |
| Anticancer        | 3ERT   | Native ligand - 4-Hydroxytamoxifen | -10                       | -                            | MET-343, THR-347, LEU-349, GLU-353, LEU-354, LEU-384, MET-388, LEU-391, ARG-394, PHE-404, GLU-419, GLY-420, ILE-424, LEU-428, GLY-521, HIS-524. | LEU-346(2), ALA-350(2), ASP-351, TRP-383, LEU-386, LEU-387, MET-421(2), LEU-525(2). |
|                   |        | Caryophyllene                      | -8.5                      | -                            | MET-343, LEU-349, TRP-383, PHE-404, ILE-424, LEU-428, GLY-521.  | LEU-346, ALA-350, LEU-384, LEU-387, MET-388, LEU-391, MET-421, LEU-525.             |
|                   |        | dihydro-cis-alpha-copaene-8-ol     | -8                        | -                            | MET-343, THR-347, TRP-383, LEU-384, ARG-394, LEU-525.   | LEU-346, ALA-350, LEU-386, LEU-387, MET-388, LEU-391, PHE-404, MET-421, LEU-428.    |
| Anti inflammatory | 2AZ5   | Native ligand - CID 5327044        | -9.2                      | GLY-121(A)                   | LEU-57(A), GLY-121(A), GLY-122(A), TYR-151(A), ILE-155(A), SER-60(B), GLN-61(B), GLY-121(B), GLY-122(B).  | TYR-59(A)(2), TYR-119(A)(2), LEU-57(B), TYR-59(B)(2), TYR-119(B)(2), TYR-151(B).    |

| Targets | PDB-ID | Compounds                      | Binding energy (kcal/mol) | H-bonds | Vander Valls interactions  | Pi interactions                           |
|---------|--------|--------------------------------|---------------------------|---------|--|---|
|         |        | Caryophyllene                  | -7                        | -       | TYR-119(A), LEU-120(A), GLY-121(A), GLY-122(A), SER-60(B), TYR-119(B), LEU-120(B), GLY-121(B), TYR-151(B), ILE-155(B). | LEU-57(A)(2), LEU-57(B)(2), TYR-59(B)(3). |
|         |        | dihydro-cis-alpha-copaene-8-ol | -7.8                      | -       | TYR-59(A), TYR-119(A), LEU-120(A), GLY-121(A), GLY-122(A), SER-60(B), GLN-61(B), LEU-120(B), GLY-121(B), TYR-151(B).   | LEU-57(B), TYR-59(B)(2), TYR-119(B)(2).   |

Bolded amino acid residues are the residues that matched with active residues of target from PDB.

## DISCUSSION

$\beta$ -pinene which was present in the leaves, roots, and stem of *H. cordata* is a well-known representative of the monoterpenes group, and is found in many plants' essential oils. There have been reports of a wide range of pharmacological activity, including the modulation of antibiotic resistance and analgesic, anticoagulant, anticancer, and antibacterial, effects that are anti-leishmanial, anti-inflammatory, anti-malarial, and antioxidants.  $\beta$ -pinene is used as an antibacterial due to its toxic effects on membranes and has been found to have inhibitory effects on leukemia and breast cancer. Leaf alcohol and leaf aldehyde are responsible for the green odor in leaves and fruits. Because of its pleasant scent and its anti-fungal and anti-inflammatory properties, Sabinene is utilized in the fragrance and flavoring industry as well as in the medicinal industry.<sup>30-32</sup>  $\alpha$ -Pinene is a monoterpene that is known to possess antimicrobial, apoptotic, anti-metastatic, and antibiotic properties.  $\alpha$ -pinene is one promising agent for the treatment of various inflammatory diseases as it has been found to suppress MAPKs and the NF- $\kappa$ B pathway.<sup>33</sup> Caryophyllene is sesquiterpenes which have anti-cancer, local anaesthetic, antimicrobial, antibacterial, anti-inflammatory, anticonvulsant, and analgesic properties.<sup>34-38</sup> Camphene is a cyclic monoterpene that has anti-viral, insecticidal, antinociceptive, and antioxidant properties.<sup>39,40</sup> o-Cymene is an insecticidal and repellent which also has antifungal and antimicrobial properties.<sup>41-43</sup> 2-Norpinene, 3,6,6-trimethyl, a chemical compound present in the leaves of *H. cordata* have antifungal properties. Due to its many benefits

and uses, including its popularity in Northeast India traditional cuisine, this plant offers positive health benefits.

## CONCLUSION

This study showed that headspace SPME-GC-MS is a very useful tool for the analysis of aromatic compounds of several edible and medicinally useful spices. *In silico* studies further supported the antimicrobial, antioxidant, anti-cancer, and anti-inflammatory properties of caryophyllene and dihydro-cis-alpha-copaene-8-ol found in *H. cordata*. Monoterpenes as the major components found in the leaves, roots, and stem of *H. cordata*, are in agreement with the *in silico* results that include antimicrobial, antioxidant, anti-inflammatories, and anti-cancer properties. Given all the beneficial active components, *H. cordata* acts as an excellent edible genetic resource and its consumption as a diet can benefit greatly in human health and promotes natural product research. Despite the potential and presence of numerous significant compounds, more research on the therapeutic applications of *H. cordata* can be performed.

## ACKNOWLEDGEMENT

We thank the Institute of Bioresources and Sustainable Development, Imphal, for both manpower and instrumentation supporting this study.

## CONFLICT OF INTEREST

The authors declared that they have no competing interests.

## ABBREVIATIONS

**VOC's:** Volatile organic compounds; **GC-MS:** Gas chromatography mass spectrometry; **SPME:** Solid phase microextraction; **NIST:** National Institute of Standards and Technology; **AM:** Anti-microbial; **AO:** Anti-oxidant; **AC:** Anti-cancer; **AI:** Anti-inflammatory; **DHFR:** Dihydrofolate reductase; **Era:** Estrogen Receptor Alpha.

## REFERENCES

- Sawian JT, Jeeva S, Lyndem FG, Mishra BP, Laloo RC. Wild edible plants of Meghalaya, North-East India. *Nat Prod Radiance*. 2007; 6(5): 410-26.
- Hynniewta SR, Kumar Y. Herbal remedies among the Khasi traditional healers and village folks in Meghalaya. *Indian J Tradit Knowl*. 2008; 7(4): 581-6.
- Deb CR, Jamir S, Sangtam TL, Jamir NS. A study on the medicinal plants used by the Sangtam Naga Tribe in Kiphire district, Nagaland, India. *J Ayurvedic Herb. Med*. 2012; 18: 2(02).
- Kala CP. Ethnomedicinal botany of the Apatani in the Eastern Himalayan region of India. *J Ethnobiol Ethnomed*. 2005; 1: 11. doi: 10.1186/1746-4269-1-11, PMID 16288657.
- Imotomba RK, Devi LS. Creation of geo-spatial data base of medicinal plants of Senapati district, Manipur. *Natl J Chembiosis*. 2011; 2(2).
- Lau KM, Lee KM, Koon CM, Cheung CS, Lau CP, Ho HM, et al. Immunomodulatory and anti-SARS activities of *Houttuynia cordata*. *J Ethnopharmacol*. 2008; 118(1): 79-85. doi: 10.1016/j.jep.2008.03.018, PMID 18479853.
- Park E, Kum S, Wang C, Park SY, Kim BS, Schuller-Levis G. Anti-inflammatory activity of herbal medicines: inhibition of nitric oxide production and tumor necrosis factor- $\alpha$  secretion in an activated macrophage-like cell line. *Am J Chin Med*. 2005; 33(3): 415-24. doi: 10.1142/S0192415X05003028, PMID 16047559.
- Lu HM, Liang YZ, Yi LZ, Wu XJ. Anti-inflammatory effect of *Houttuynia cordata* injection. *J Ethnopharmacol*. 2006; 104(1-2): 245-9. doi: 10.1016/j.jep.2005.09.012, PMID 16213118.
- Li GZ, Chai OH, Lee MS, Han EH, Kim HT, Song CH. Inhibitory effects of *Houttuynia cordata* water extracts on anaphylactic reaction and mast cell activation. *Biol Pharm Bull*. 2005; 28(10): 1864-8. doi: 10.1248/bpb.28.1864, PMID 16204936.
- Hayashi K, Kamiya M, Hayashi T. Virucidal effects of the steam distillate from *Houttuynia cordata* and its components on HSV-1, influenza virus, and HIV. *Planta Med*. 1995; 61(3): 237-41. doi: 10.1055/s-2006-958063, PMID 7617766.
- Cheng A, Zhang W, Xie Y, Jiang W, Arnold E, Sarafianos SG, et al. Expression, purification, and characterization of SARS coronavirus RNA polymerase. *Virology*. 2005; 335(2): 165-76. doi: 10.1016/j.virol.2005.02.017, PMID 15840516.
- Kim SK, Ryu SY, Choi SU, Kim YS. Cytotoxic alkaloids from *Houttuynia cordata*. *Arch Pharm Res*. 2001; 24(6): 518-21. doi: 10.1007/BF02975156, PMID 11794526.
- Singh NB, Devi ML, Biona T, Sharma N, Das S, Chakravorty J, et al. Phytochemical composition and antimicrobial activity of essential oil from the leaves of *Artemisia vulgaris* L. *Molecules*. 2023 Feb 28; 28(5): 2279. doi: 10.3390/molecules28052279, PMID 36903525.
- Deng H, He R, Huang R, Pang C, Ma Y, Xia H, et al. Optimization of a static headspace GC-MS method and its application in metabolic fingerprinting of the leaf volatiles of 42 citrus cultivars. *Front Plant Sci*. 2022; 13: 1050289. doi: 10.3389/fpls.2022.1050289, PMID 36570894.
- Linstrom PJ, Mallard WG. The NIST Chemistry WebBook: A chemical data resource on the internet. *J Chem Eng Data*. 2001; 46(5): 1059-63. doi: 10.1021/je000236i.
- Kim S, Chen J, Cheng T, Gindulyte A, He J, He S, et al. PubChem in 2021: new data content and improved web interfaces. *Nucleic Acids Res*. 2021; 49: 1388-95.
- Morris GM, Huey R, Lindstrom W, Sanner MF, Belew RK, Goodsell DS, et al. AutoDock4 and AutoDockTools4: automated docking with selective receptor flexibility. *J Comput Chem*. 2009; 30(16): 2785-91. doi: 10.1002/jcc.21256, PMID 19399780.
- Petersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, et al. UCSF Chimera—a visualization system for exploratory research and analysis. *J Comput Chem*. 2004; 25(13): 1605-12. doi: 10.1002/jcc.20084, PMID 15264254.
- O'Boyle NM, Banck M, James CA, Morley C, Vandermeersch T, Hutchison GR. Open Babel: an open chemical toolbox. *J Cheminform*. 2011; 3: 33. doi: 10.1186/1758-2946-3-33, PMID 21982300.
- Jianu C, Stoin D, Cocan I, David I, Pop G, Lukinich-Gruia AT, et al. *In silico* and *in vitro* evaluation of the antimicrobial and antioxidant potential of *Mentha x smithiana* R. Graham Essential Oil from Western Romania. *Foods*. 2021; 10(4): 815. doi: 10.3390/foods10040815, PMID 33918674.
- Audat SA, Al-Shar'i NA, Al-Oudat BA, Bryant-Friedrich A, Bedi MF, Zayed AL, Al-Balas QA. Identification of human leukotriene A4 hydrolase inhibitors using structure-based pharmacophore modeling and molecular docking. *Molecules*. 2020; 25: 2871.
- Trott O, Olson AJ. AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *J Comput Chem*. 2010; 31(2): 455-61. doi: 10.1002/jcc.21334, PMID 19499576.
- Schrödinger L, DeLano W. PyMOL; 2020. Available from: <http://www.pymol.org/py/mol>.
- Reeve SM, Si D, Krucinska J, Yan Y, Viswanathan K, Wang S, et al. Toward broad spectrum dihydrofolate reductase inhibitors targeting trimethoprim resistant enzymes identified in clinical isolates of methicillin resistant *Staphylococcus aureus*. *ACS Infect Dis*. 2019; 5(11): 1896-906. doi: 10.1021/acinfecdis.9b00222, PMID 31565920.
- Bury-Moné S. Antibacterial therapeutic agents: antibiotics and bacteriophages.
- Raimondi MV, Randazzo O, La Franca M, Barone G, Vignoni E, Rossi D, et al. DHFR inhibitors: reading the past for discovering novel anticancer agents. *Molecules*. 2019; 24(6): 1140. doi: 10.3390/molecules24061140, PMID 30909399.
- He L, He T, Farrar S, Ji L, Liu T, Ma X. Antioxidants maintain cellular redox homeostasis by elimination of reactive oxygen species. *Cell Physiol Biochem*. 2017; 44(2): 532-53. doi: 10.1159/000485089, PMID 29145191.
- Costa JDS, Ramos RDS, Costa KDSL, Brasil DDSB, Silva CHTPD, Ferreira EFB, et al. An *in silico* study of the antioxidant ability for two caffeine analogs using molecular docking and quantum chemical methods. *Molecules*. 2018; 23(11): 2801. doi: 10.3390/molecules23112801, PMID 30380600.
- Muchtaridi M, Syahidah HN, Subarnas A, Yusuf M, Bryant SD, Langer T. Molecular docking and 3D-pharmacophore modeling to study the interactions of chalcone derivatives with estrogen receptor alpha. *Pharmaceuticals (Basel)*. 2017; 10(4): 81. doi: 10.3390/ph10040081, PMID 29035298.
- Abdullahi SH, Uzairu A, Shallangwa GA, Uba Sani, Umar AB. Structure based design of some novel 3-methylquinoxaline derivatives through molecular docking and pharmacokinetics studies as novel VEGFR-2 inhibitors. *Chem Afr*. 2022; 5(6): 1967-78. doi: 10.1007/s42250-022-00485-3.
- Gerriets V, Goyal A, Khaddour K. Tumor necrosis factor inhibitors. *InStatPearls*. StatPearls Publishing. 2021.18.
- Lis K, Kuzawi ska O, Bałkowiec-Iskra E. Tumor necrosis factor inhibitors: state of knowledge. *Arch Med Sci*. 2014; 10: 1175-85.
- Zia K, Ashraf S, Jabeen A, Saeed M, Nur-e-Alam M, Ahmed S, et al. Identification of potential TNF- $\alpha$  inhibitors: from *in silico* to *in vitro* studies. *Sci Rep*. 2020; 10(1): 20974. doi: 10.1038/s41598-020-77750-3, PMID 33262408.
- Cao Y, Zhang H, Liu H, Liu W, Zhang R, Xian M, et al. Biosynthesis and production of sabinene: current state and perspectives. *Appl Microbiol Biotechnol*. 2018; 102(4): 1535-44. doi: 10.1007/s00253-017-8695-5, PMID 29264773.
- Menon AN, Padmakumari KP. Studies on essential oil composition of cultivars of black pepper (*Piper nigrum* L.)—V. *J Essent Oil Res*. 2005; 17(2): 153-5. doi: 10.1080/10412905.2005.9698862.
- Valente J, Zuzarte M, Gonçalves MJ, Lopes MC, Cavaleiro C, Salgueiro L, et al. Antifungal, antioxidant and anti-inflammatory activities of *Oenanthe crocata* L. essential oil. *Food Chem Toxicol*. 2013; 62: 349-54. doi: 10.1016/j.fct.2013.08.083, PMID 24012643.
- Kim DS, Lee HJ, Jeon YD, Han YH, Kee JY, Kim HJ, et al. Alpha-pinene exhibits anti-inflammatory activity through the suppression of MAPKs and the NF- $\kappa$ B pathway in mouse peritoneal macrophages. *Food Chem Toxicol*. 2015; 43: 731-42.
- Legault J, Côté PA, Simard S, Pichette A. Iso-caryophyllene cytotoxicity induced by lipid peroxidation and membrane permeabilization in L-929 cells. *J Appl Pharm Sci*. 2013; 3: 025-31.
- Ghelardini C, Galeotti N, Di Cesare Mannelli LD, Mazzanti G, Bartolini A. Local anaesthetic activity of  $\beta$ -caryophyllene. *Farmaco*. 2001; 56(5-7): 387-9. doi: 10.1016/S0014-827X(01)01092-8, PMID 11482764.
- Chavan MJ, Wakte PS, Shinde DB. Analgesic and anti-inflammatory activity of caryophyllene oxide from *Annona squamosa* L. bark. *Phytomedicine*. 2010; 17(2): 149-51. doi: 10.1016/j.phymed.2009.05.016, PMID 19576741.
- Fidyk K, Fiedorowicz A, Strzdała L, Szumny A.  $\beta$ -caryophyllene and  $\beta$ -caryophyllene oxide—natural compounds of anticancer and analgesic properties. *Cancer Med*. 2016; 5(10): 3007-17. doi: 10.1002/cam4.816, PMID 27696789.
- Sokolova AS, Putilova VP, Yarovaya OI, Zybikina AV, Mordvinova ED, Zaykovskaya AV, et al. Synthesis and antiviral activity of camphene derivatives against different types of viruses. *Molecules*. 2021; 26(8): 2235. doi: 10.3390/molecules26082235, PMID 33924393.
- Pavela R, Maggi F, Mazzara E, Torresi J, Cianfaglione K, Benelli G, et al. Prolonged sublethal effects of essential oils from non-wood parts of nine conifers on key insect pests and vectors. *Ind Crops Prod*. 2021; 168: 113590. doi: 10.1016/j.indcrop.2021.113590.

**Cite this article:** Biona T, Singh NB, Keithellakpam OS, Srinivasan S, Patil S, Mukherjee PK. Headspace SPME-GC-MS Analysis and *in silico* Molecular Docking Studies of Phytochemical Compounds Present in *Houttuynia cordata* Thunb. *Indian J of Pharmaceutical Education and Research*. 2024;58(3):870-81.